



# ANNALES DE L'INSTITUT FOURIER

Farshid HAJIR, Christian MAIRE & Ravi RAMAKRISHNA

**Deficiency of  $p$ -class tower groups and Minkowski units**

Article à paraître, mis en ligne le 5 février 2025, 48 p.

Article mis à disposition par ses auteurs selon les termes de la licence  
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE



<http://creativecommons.org/licenses/by-nd/3.0/fr/>



Les *Annales de l'Institut Fourier* sont membres du  
Centre Mersenne pour l'édition scientifique ouverte

[www.centre-mersenne.org](http://www.centre-mersenne.org)

e-ISSN : 1777-5310

## DEFICIENCY OF $p$ -CLASS TOWER GROUPS AND MINKOWSKI UNITS

by Farshid HAJIR,  
Christian MAIRE & Ravi RAMAKRISHNA (\*)

---

ABSTRACT. — The deficiency  $\text{Def}(G)$  of a finitely presented pro- $p$  group  $G$  is the difference between its minimal numbers of relations and generators. For a number field  $K$  with maximal unramified  $p$ -extension  $K_\theta$ , set  $G_\theta = \text{Gal}(K_\theta/K)$ . Shafarevich (and independently Koch) showed

$$0 \leq \text{Def}(G_\theta) \leq \dim(\mathcal{O}_K^\times / (\mathcal{O}_K^\times)^p).$$

We explore connections between the relations of  $G_\theta$  and the Galois module structure of the units in the tower  $K_\theta/K$ . If  $\mu_p \not\subset K$ , we give an exact formula for  $\text{Def}(G_\theta)$  in terms of the number of independent Minkowski units in the tower. We also study the depth of the relations of  $G_\theta$  in the Zassenhaus filtration and provide evidence that the Shafarevich–Koch upper bound is “almost always” sharp. In the other direction, we give the first examples of infinite  $G_\theta$  with  $\text{Def}(G_\theta) = 0$  and  $\dim(\mathcal{O}_K^\times / (\mathcal{O}_K^\times)^p)$  large, so that the upper bound is not sharp.

RÉSUMÉ. — Le défaut  $\text{Def}(G)$  d’un pro- $p$  groupe  $G$  de type fini est la différence entre son nombre minimal de relations et son nombre minimal de générateurs. Pour un corps de nombres  $K$  de  $p$ -extension non ramifiée maximale  $K_\theta$ , posons  $G_\theta = \text{Gal}(K_\theta/K)$ . Shafarevich (et indépendamment Koch) a montré que

$$0 \leq \text{Def}(G_\theta) \leq \dim(\mathcal{O}_K^\times / (\mathcal{O}_K^\times)^p).$$

Nous explorons les connexions entre les relations de  $G_\theta$  et la structure galoisienne des unités dans la tour  $K_\theta/K$ . Si  $\mu_p \not\subset K$ , nous donnons une formule exacte pour

---

*Keywords:* Deficiency, Golod–Shafarevich polynomial,  $p$ -class field tower, Zassenhaus filtration.

2020 *Mathematics Subject Classification:* 11R29, 11R37.

(\*) This work started when the second author held a visiting scholar position at Cornell University, funded by the program “Mobilité sortante” of the Région Bourgogne Franche-Comté, during the 2017-18 academic year. It continued during visits to the Harbin Institute of Technology and Cornell University. CM thanks the Department of Mathematics at Cornell University and the Institute for Advanced Study in Mathematics of HIT for providing excellent conditions for conducting research. The second author was partially supported by the ANR project FLAIR (ANR-17-CE40-0012) and by the EIPHI Graduate School (ANR-17-EURE-0002). The third author was supported by Simons Collaboration grant #524863. All three authors were supported by Mathematisches Forschungsinstitut Oberwolfach for a Research in Pairs visit in January, 2019 and by ICERM for a Research in Pairs visit in January, 2020.

$\text{Def}(G_\emptyset)$  en termes du nombre d'unités de Minkowski indépendantes dans la tour. Nous étudions également la profondeur des relations de  $G_\emptyset$  suivant la filtration de Zassenhaus et montrons que la borne supérieure de Shafarevich–Koch est « presque toujours » optimale. Mais dans l'autre sens, nous donnons les premiers exemples pour lesquels  $G_\emptyset$  est infini avec  $\text{Def}(G_\emptyset) = 0$  et  $\dim(\mathcal{O}_K^\times/(\mathcal{O}_K^\times)^p)$  grand, de sorte que la borne supérieure n'est pas optimale.

## Introduction

Let  $p$  be a prime number, and let  $K$  be a number field. For a finite set  $S$  of places of  $K$ , let  $K_S$  be the maximal  $p$ -extension of  $K$  unramified outside  $S$  and  $G_S = \text{Gal}(K_S/K)$ , its Galois group. Note in particular that  $K_\emptyset$  is the maximal pro- $p$  extension of  $K$  unramified everywhere; we call  $K_\emptyset/K$  the  $p$ -class field tower of  $K$ . Let

$$d(G_S) = \dim H^1(G_S) = \dim H^1(G_S, \mathbb{Z}/p)$$

$$\text{and } r(G_S) = \dim H^2(G_S) = \dim H^2(G_S, \mathbb{Z}/p)$$

be, respectively, the minimal number of generators and relations of  $G_S$ . Define  $\text{Def}(G_S) := r(G_S) - d(G_S)$  as the *deficiency* of  $G_S$ .<sup>(1)</sup>

Our goal in this paper is to better understand the pro- $p$  group  $G_S$  when  $S$  is tame, that is when  $S$  does not contain any places above  $\{p, \infty\}$ . We are particularly interested in the case  $S = \emptyset$ . In the wild setting, when  $S$  contains all places above  $\{p, \infty\}$ , we have at our disposal a very powerful tool, namely the global duality theorem for  $G_S$ , which immediately yields an explicit and easily computable formula for  $\text{Def}(G_S)$ . While such a tool is still missing in the tame, and in particular, unramified, case, Theorem A below (see Theorem 2.9) is a step toward refining our understanding of the unramified situation by relating it to the presence of Minkowski units in  $K_\emptyset/K$ .

By class field theory, the maximal abelian quotient of  $G_S$  is isomorphic to the  $p$ -Sylow subgroup of a ray class group for  $K$ , and is therefore finite when  $S$  is tame. By the Burnside Basis Theorem,  $d(G_\emptyset)$  is the  $p$ -rank of the class group of  $K$  and is in particular computable in any given case (at least in theory). In contrast, we do not know an algorithm for computing  $r(G_\emptyset)$ . However, thanks to the celebrated work of Shafarevich [37] (and, independently, Koch – see for example [15, Chapter 11]) we know that

$$0 \leq \text{Def}(G_\emptyset) \leq d(\mathcal{O}_K^\times),$$

---

<sup>(1)</sup>In most of the group theory literature the deficiency of  $G$  is defined as  $d(G) - r(G)$ .

where  $d(\mathcal{O}_K^\times) := d(\mathcal{O}_K^\times/(\mathcal{O}_K^\times)^p)$  is the  $p$ -rank of the unit group  $\mathcal{O}_K^\times$  of the ring of integers  $\mathcal{O}_K$  of  $K$ . We recall that if  $K$  has  $r_1$  embeddings into  $\mathbb{R}$  and  $r_2$  pairs of complex conjugate embeddings into  $\mathbb{C}$ , then  $d(\mathcal{O}_K^\times) = r_1 + r_2 - 1 + \delta$  where  $\delta$  is 1 or 0 according to whether  $K$  contains a primitive  $p$ th root of unity  $\zeta_p$  or not. The number-theoretic work of Shafarevich on the above deficiency bound [37] subsequently led to the group-theoretic work of Golod and Shafarevich [9]. This pair of papers gave a criterion for the infinitude of  $G_\emptyset$  and produced the first such examples. More historical details can be found in [15, 24, 28, 36].

In this work, we investigate more closely the relationship between units in unramified class field towers and defining relations for their Galois groups. Our main theorem is that the existence of certain types of units along the tower  $K_\emptyset/K$  provides two types of results: (a) tighter bounds for the deficiency of  $G_\emptyset$ , as well as (b) more refined information on the depth of its defining relations.

We introduce a constant  $\lambda$  measuring the free-part of the Galois module structure of the units in  $K_\emptyset/K$  (see Section 2.3). Here free-part means the following: if  $F/K$  is a finite Galois extension in  $K_\emptyset/K$  with Galois group  $G$ , we are interested in the  $\mathbb{F}_p[G]$ -structure of  $\mathbb{F}_p \otimes \mathcal{O}_F^\times$ , that is the units  $\mathcal{O}_F^\times$  modulo  $p$ th powers. Recall that since  $G$  is a  $p$ -group, the category of  $\mathbb{F}_p[G]$ -modules is not semisimple. When the  $\mathbb{F}_p[G]$ -free part of  $\mathbb{F}_p \otimes \mathcal{O}_F^\times$  is nontrivial, we say the extension  $F/K$  admits a *Minkowski unit* (see Section 1.3 for further details). It is not difficult to see that the number of independent Minkowski units is non-increasing and stabilizes as we move up the tower  $K_\emptyset/K$  and therefore after a finite number of steps reaches a constant value we denote  $\lambda := \lambda_{K_\emptyset/K}$ . We also define  $\beta$  to be

$$\beta := \begin{cases} d\left(\frac{\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\emptyset}^\times)^p}{(\mathcal{O}_K^\times)^p}\right) & \zeta_p \in K \\ 0 & \text{otherwise.} \end{cases}$$

Note that when  $\zeta_p \in K$ , if we set  $L = K_\emptyset \cap K(\sqrt[p]{\mathcal{O}_K^\times})$ , then  $[L : K] = p^\beta$ . Thus,  $0 \leq \beta \leq \min(r_1 + r_2, d(G_\emptyset))$ . We also note that  $\beta > 0$  if and only if for some  $u \in \mathcal{O}_K^\times$ ,  $K(u^{1/p})/K$  is a  $\mathbb{Z}/p$ -unramified extension. More generally, the quantity  $\beta$  quantifies the number of such independent extensions of  $K$ .

**THEOREM A.** — *With notations as above, for the deficiency of the  $p$ -class tower group  $G_\emptyset$ , we have the following upper and lower bounds:*

$$d(\mathcal{O}_K^\times) - \lambda - \beta \leq \text{Def}(G_\emptyset) \leq d(\mathcal{O}_K^\times) - \lambda.$$

In particular if  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\theta}^\times)^p = (\mathcal{O}_K^\times)^p$  or if  $K$  does not contain a primitive  $p$ th root of unity, then  $\text{Def}(G_\theta) = d(\mathcal{O}_K^\times) - \lambda$ .

We give two proofs of this result (see proof of Theorem 2.9). In our second more constructive proof we realize  $G_\theta$  as a quotient of some  $G_S$ , where  $S$  is a well-chosen finite set of tame (coprime to  $p$ ) prime ideals of  $\mathcal{O}_K$  (see also Notations at the end of this section), and we use the Hochschild–Serre spectral sequence induced by the natural map  $G_S \twoheadrightarrow G_\theta$  to produce  $d(G_\theta) + d(\mathcal{O}_K^\times) - \lambda - \beta$  independent elements of  $\text{III}_\theta^2 = H^2(G_\theta)$ .

In Section 3.6 we also give examples of infinite  $G_\theta$  with as many as seven independent Minkowski units.

*Remark.* — As mentioned earlier, the non-negativity of  $\text{Def}(G_\theta)$  follows from a basic group-theoretical property of  $G_\theta$ , namely that its maximal abelian quotient is finite. In other words, one knows that  $G_\theta$  has at least  $d(G_\theta)$  relations. For the group  $G_\theta$ , one can concretely produce  $d(G_\theta)$  relations for  $G_\theta$  as follows. In Figure 1, we show a tower of fields  $K \subset K' \subset L'_1 \subset L'_2$  for whose definition, the reader may consult the Notations section at the end of this introduction. For the moment, the key point is that  $L'_2/L'_1$  is an elementary abelian  $p$ -extension of dimension  $d(G_\theta)$ . By the Chebotarev density theorem, we can choose  $d(G_\theta)$  primes of  $K$  whose Frobenius automorphisms form a basis of the elementary  $p$ -abelian group  $\text{Gal}(L'_2/L'_1)$ . Letting  $S_1$  be the set consisting of these primes, in Section 2.2, we will describe in detail how applying the Gras–Munnier Theorem (Theorem 1.1) and Lemma 1.5 (ii) to the primes in  $S_1$  gives us  $d(G_\theta)$  independent elements in  $\text{III}_\theta^2 = H^2(G_\theta)$ , which in turn correspond to  $d(G_\theta)$  distinct relations in a minimal presentation of  $G_\theta$ . We refer to relations constructed in this way as “easily detected” via  $S_1$ , or, when the context is clear, simply “easy.”

A key observation we make in this work is that aside from the  $d(G_\theta)$  relations easily detected via  $S_1$ , we can construct  $d(\mathcal{O}_K^\times) - \lambda - \beta$  additional relations via a modification of this construction using a further set  $S_2$  of auxiliary primes whose Frobenius automorphisms span a Galois group in a more complicated tower of governing fields (Figure 2.1) described in Section 2.2. The existence of such primes is tied up with the Galois module structure of units in the Hilbert  $p$ -class field tower. We refer to the resulting relations as “difficult” relations “detected” by  $S_2$ . This set of ideas leads to the lower bound for  $\text{Def}(G_\theta)$  in the theorem. The upper bound, on the other hand, is a consequence of a result of Wingberg [40]. When  $\beta = 0$  (which is always the case if  $K$  does not contain a primitive  $p$ th root of unity), these upper and lower bounds coincide, in which case all the relations are either




Figure 0.1.

easily detected by  $S_1$  or difficult and detected by  $S_2$ . But when  $\beta > 0$ , only  $d(\mathcal{O}_K^\times) - \lambda - \beta$  of the relations are constructible in this way. Indeed, in the example below  $\beta = 1$  and we find the final relation, which is difficult, by an ad hoc method.

*Example.* — Take  $p = 2$ . Our method allows us to show that for  $K = \mathbb{Q}(\sqrt{-5460})$ , the example studied extensively by Boston–Wang [4], there are 4 easily detected relations and another relation that is difficult.  $\text{Def}(G_\emptyset) = r - d = 5 - 4 = 1$ .

Thanks to the work of Labute [16], Schmidt [31] and others we know that there are special sets  $S$  (finite and tame) for which  $G_S$  is of cohomological dimension 2; however, their methods do not allow  $S$  to be empty. In particular, the question of the computation of the cohomological dimension of  $G_\emptyset$  has only been resolved in a few cases, namely when  $G_\emptyset$  is known to be finite. A consequence of Theorem A is the following (Theorem 3.12):

COROLLARY. — *Let  $K$  be a number field such that*

- (i)  $K$  contains a primitive  $p$ th root of unity;
- (ii)  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_0}^\times)^p = (\mathcal{O}_K^\times)^p$ .

Then  $\dim H^3(G_\emptyset, \mathbb{F}_p) > 0$ . Moreover:

- If  $\dim H^3(G_\emptyset, \mathbb{F}_p) = 1$ , then  $G_\emptyset$  is finite or of cohomological dimension 3;
- If  $\text{Def}(G_\emptyset) = 0$  and  $G_\emptyset$  is of cohomological dimension 3, then  $G_\emptyset$  is a Poincaré duality group.

We also deduce (Corollary 3.4):

COROLLARY. — *Let  $K$  be a number field such that*

- (i)  $K$  contains a primitive  $p$ th root of unity;
- (ii)  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\theta}^\times)^p = (\mathcal{O}_K^\times)^p$ ;
- (iii)  $\text{Def}(G_\theta) = 0$ .

*Then for every open subgroup  $H$  of  $G_\theta$ , one has  $\text{Def}(H) = 0$ .*

When  $\text{Def}(H) = 0$  for all open subgroups of  $G$ ,  $\lambda$  is maximal all along the tower. When  $\text{Def}(G_\theta) = 0$  and the tower is finite, it is a standard result from finite group theory that  $G_\theta$  is either cyclic or, when  $p = 2$ , a generalized quaternion group. Observe also that Poincaré groups of dimension 3 have deficiency zero.

When  $G_\theta$  is infinite we suspect  $\text{Def}(G_\theta)$  is maximal (namely equal to  $d(\mathcal{O}_K^\times)$ ) very often in accordance with the heuristics of Liu, Wood and Zureick-Brown [19]. In fact, we elaborate a strategy to investigate maximality of the deficiency by testing for the presence of Minkowski units through computer computation. We further note that if in the first steps of the tower  $K_\theta/K$  there are  $\alpha$  Minkowski units preventing us from concluding that  $\text{Def}(G_\theta)$  is maximal, the group  $G_\theta$  can be described by at least  $\alpha$  relations of high depth in the Zassenhaus filtration. Denote by  $(K_n)$  the sequence in  $K_\theta/K$  where  $K_1 := K$  and  $K_{n+1}$  is the maximal elementary abelian  $p$ -extension of  $K_n$  in  $K_\theta/K$ . Put  $H_n = \text{Gal}(K_n/K)$ . Let  $r_{\max} = d(G_\theta) + d(\mathcal{O}_K^\times)$  be the maximal possible value of  $r(G_\theta)$ . To each presentation of a pro- $p$  group, there is associated a Golod–Shafarevich polynomial; for the basic facts of these polynomials, see Section 4.1.2. Golod and Shafarevich proved that if this polynomial vanishes on the open unit interval, then the group must be infinite. In Section 4, we prove the following result (see Theorem 4.12). See also Section 1.3.3.

THEOREM B. — *Let  $\lambda_n$  be the number of independent  $\mathbb{F}_p[H_n]$ -Minkowski units in  $K_n$ . Then  $G_\theta$  can be generated by  $d(G_\theta)$  generators and  $r_{\max}$  relations  $\{\rho_1, \dots, \rho_{r_{\max}}\}$  such that at least  $\lambda_n$  relations are of depth greater than  $2^n$ . Hence, we can take  $1 - d(G_\theta)t + (r_{\max} - \lambda_n)t^2 + \lambda_n t^{2^n}$  as a Golod–Shafarevich polynomial for  $G_\theta$ .*

The more familiar Golod–Shafarevich polynomial in this context is  $1 - d(G_\theta)t + r_{\max}t^2$ , which is less likely to have a root and thus indicate  $\#G_\theta = \infty$ . Also, we will allow the possibility of  $n = \infty$ , that is there may be fewer than  $r_{\max}$  relations.

The imaginary quadratic case is particularly easy to study (here  $p = 2$ ). Indeed when  $K$  is an imaginary quadratic field, one has  $\text{Def}(G_\theta) \in \{0, 1\}$ . Here we show that, almost always, there is no Minkowski unit in any

quadratic extension  $F/K$  in  $K_\emptyset/K$ , which implies  $\text{Def}(G_\emptyset) = 1$  (see Theorem 5.12). Denote by  $\mathcal{F}$  the set of imaginary quadratic fields, and for  $X \geq 2$ , put

$$\mathcal{F}(X) = \{K \in \mathcal{F}, |\text{disc}(K)| \leq X\}, \quad \mathcal{F}_0(X) = \{K \in \mathcal{F}(X), \text{Def}(G_\emptyset) = 0\}.$$

THEOREM C. — *Let  $p = 2$ . One has*

$$\frac{\#\mathcal{F}_0(X)}{\#\mathcal{F}(X)} \leq C \frac{\log \log X}{\sqrt{\log X}},$$

where  $C$  is an absolute constant and  $X$  is large enough. In particular, the proportion of imaginary quadratic fields of discriminant at most  $X$  for which  $\text{Def}(G_\emptyset) = 0$  tends to zero as  $X \rightarrow \infty$ .

Notations.

- Let  $p$  be a prime number and  $K$  be a number field.
- We denote by
  - $\mathcal{O}_K$  the ring of integers of  $K$ , and by  $\mathcal{O}_K^\times$  the group of units of  $\mathcal{O}_K$ ,
  - $\mathcal{E}_K = \mathbb{F}_p \otimes \mathcal{O}_K^\times$ , the units modulo the  $p$ th-powers,
  - $K^H$  the Hilbert  $p$ -class field of  $K$ ,
  - $\text{Cl}_K$  the  $p$ -Sylow subgroup of class group of  $K$ .
- Let  $\zeta_p \in \mathbb{Q}^{alg}$  be a primitive  $p$ th root of 1. Put  $\delta := \delta_{K,p} := 1$  when  $\zeta_p \in K$ , 0 otherwise.
- Let  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_s\}$  be a finite set of prime ideals of  $K$  of cardinality  $s$ . We identify a prime  $\mathfrak{p} \in S$  with the place  $v$  it defines.
  - We assume each  $\mathfrak{p}_i$  is tame (prime to  $p$ ) and satisfies  $|\mathcal{O}_K/\mathfrak{p}_i| \equiv 1 \pmod{p}$ .
  - We denote by  $\text{RCG}_K(\mathfrak{p}_1, \dots, \mathfrak{p}_s)$  the  $p$ -Sylow subgroup of the ray class group of  $K$  of modulus  $\mathfrak{p}_1 \cdots \mathfrak{p}_s$ . When  $S = \emptyset$ , one has  $\text{RCG}_K(\emptyset) = \text{Cl}_K$ .
  - Let  $K_S$  be the maximal pro- $p$  extension of  $K$  unramified outside  $S$ , put  $G_S = G_{K,S} = \text{Gal}(K_S/K)$ .
  - By class field theory, one has  $G_S^{ab} \simeq \text{RCG}_K(\mathfrak{p}_1, \dots, \mathfrak{p}_s)$ .
  - Put  $V_S := \{x \in K^\times, (x) = I^p \text{ as a fractional ideal of } K; x \in (K_v^\times)^p, \forall v \in S\}$ . Then  $V_S \supset (K^\times)^p$  and we have the exact sequence:

$$0 \longrightarrow \mathcal{O}_K^\times / \mathcal{O}_K^{\times p} \longrightarrow V_\emptyset / (K^\times)^p \longrightarrow \text{Cl}_K[p] \longrightarrow 0.$$

The group  $V_\emptyset / (K^\times)^p$  is often called the *Selmer group* of  $K$ .

- If  $M$  is a  $\mathbb{Z}$ -module, we set  $d(M) = \dim_{\mathbb{F}_p}(\mathbb{F}_p \otimes M)$ .



- When  $G$  is a pro- $p$  group, we denote  $d(G) = d(G^{ab})$ , where  $G^{ab} = G/[G, G]$ .
- Let  $(r_1, r_2)$  be the signature of  $K$ . By Dirichlet's Theorem  $d(\mathcal{O}_K^\times) = r_1 + r_2 - 1 + \delta$ .
- From the exact sequence above,  $d(V_\emptyset/(K^\times)^p) = d(\mathcal{O}_K^\times) + d(Cl_K)$ .
- Unless otherwise specified, all cohomology groups have  $\mathbb{Z}/p$ -coefficients.
  - Hence  $d(G_\emptyset) := d(H^1(G_\emptyset)) = \dim H^1(G_\emptyset, \mathbb{Z}/p)$  and  $r(G_\emptyset) := \dim H^2(G_\emptyset, \mathbb{Z}/p)$ .
  - The deficiency  $\text{Def}(G_\emptyset)$  of  $G_\emptyset$  is defined to be  $r(G_\emptyset) - d(G_\emptyset)$ .

For the computations in this paper we have used the programs GP-PARI [26] and MAGMA [3] and have often assumed the GRH to speed up the computations.

## 1. Preliminaries

In this section we develop the results we need to detect elements of  $H^2(G_\emptyset) = \text{III}_\emptyset^2$  as described in the Remark in the Introduction. In particular, Lemma 1.5 shows how one can detect elements of  $\text{III}_\emptyset^2$  via ramified extensions of  $K_\emptyset$ ; we illustrate our strategy by finding  $\text{III}_\emptyset^2$  for the field  $\mathbb{Q}(\sqrt{-5460})$  with  $p = 2$ .

In Section 1.2, we relate  $\text{Def}(G_\emptyset)$  to norms of units from number fields in the tower  $K_\emptyset/K$ . In Section 1.3 we develop the basics of the theory of Minkowski units and show, using the Gras–Munnier Theorem 1.1, that the existence of a Minkowski unit in some number field  $F$  in the tower  $K_\emptyset/K$  follows when  $G_{F, \{\mathfrak{p}\}}^{ab} = G_{F, \emptyset}^{ab}$  for some prime  $\mathfrak{p}$  of  $K$ .

### 1.1. Saturated sets, and a spectral sequence

#### 1.1.1. Degree- $p$ cyclic extensions with prescribed ramification

Take  $p$ ,  $K$  and  $S$  as in the “Notations”. The fields  $L'_i$  of Figure 0.1 are called *governing fields* as the existence of a  $\mathbb{Z}/p$ -extension of  $K$  ramified exactly at a given set of primes depends on their Frobenius automorphisms in these extensions. See Theorem 1.1 below.

For each prime ideal  $\mathfrak{p} \in S$ , let us choose a prime ideal  $\mathfrak{P}|\mathfrak{p}$  of  $\mathcal{O}_{L'_2}$ , and denote by  $\sigma_{\mathfrak{p}} := \left( \frac{L'_2/K'}{\mathfrak{P}} \right)$ , the Frobenius at  $\mathfrak{P}$  in the governing extension  $L'_2/K'$ .

Using that  $L'_2$  is formed by taking  $p$ th roots of elements of  $K$  (not  $K'$ ), one can easily show that  $\sigma_{\mathfrak{p}}$  depends, up to a nonzero scalar multiple, only on  $\mathfrak{p}$ . This serves our purposes. By abuse we also denote by  $\sigma_{\mathfrak{p}}$  its restriction to  $L'_1$ . One says that the Frobenius automorphisms  $\sigma_{\mathfrak{p}}$ ,  $\mathfrak{p} \in S$ , satisfy a *nontrivial relation* if

$$\prod_{\mathfrak{p} \in S} \sigma_{\mathfrak{p}}^{a_{\mathfrak{p}}} = 1,$$

in  $\text{Gal}(L'_2/K')$  (or in  $\text{Gal}(L'_1/K')$ ) with the  $a_i \in \mathbb{Z}/p$  not all zero. Thus the existence of a nontrivial relation is independent of the ambiguity in the choice of  $\sigma_{\mathfrak{p}}$ .

**THEOREM 1.1** (Gras–Munnier [12]). — *Let  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t\}$  be a set of tame prime ideals of  $K$ . One has:*

- (i)  $d(G_S) \neq d(G_{\emptyset})$ , if and only if the  $\sigma_{\mathfrak{p}}$ ,  $\mathfrak{p} \in S$ , satisfy a nontrivial relation in  $\text{Gal}(L'_2/K')$ .
- (ii)  $|G_S^{ab}| > |G_{\emptyset}^{ab}|$  if and only if the  $\sigma_{\mathfrak{p}}$ ,  $\mathfrak{p} \in S$ , satisfy a nontrivial relation in  $\text{Gal}(L'_1/K')$ .

For a generalization of Theorem 1.1, see [10, Chapter V].

*Remark 1.2.*

- (1) The distinction between (i) and (ii) of Theorem 1.1 is that (ii) can occur when  $d(G_S) = d(G_{\emptyset})$ . The governing fields for the width and depth of  $G_S^{ab}$  are different.
- (2) The Kummer radical of  $L'_1/K'$  is  $\mathcal{O}_K^{\times}(K'^{\times})^p/(K'^{\times})^p$  which is isomorphic to  $\mathcal{E}_K$  since  $\mathcal{O}_K^{\times} \cap (K'^{\times})^p = (\mathcal{O}_K^{\times})^p$  and  $[K' : K]$  is coprime to  $p$ . For the same reason,  $V_{\emptyset}/(K^{\times})^p$  is the Kummer radical of  $L'_2/K'$ .

### 1.1.2. Saturated sets

For  $v \in S$ , we denote by  $G_v$  the absolute Galois group of the maximal pro- $p$  extension  $\overline{K}_v$  of the completion  $K_v$  of  $K$  at  $v$ . Let  $\text{III}_S^2$  be the kernel of the localization map of  $H^2(G_S)$ :

$$\text{III}_S^2 := \ker(H^2(G_S) \longrightarrow \oplus_{v \in S} H^2(G_v)).$$

Put  $\mathbb{B}_S = (V_S/(K^{\times})^p)^{\vee}$ ; then one has (see Theorem 11.3 of [15]) a very important natural injective map  $\text{III}_S^2 \hookrightarrow \mathbb{B}_S$ . When  $S$  contains the places of  $K$  above  $\{p, \infty\}$  this map is an isomorphism and  $\text{III}_S^2$  is dual to

$$\text{III}_S^1(\mu_p) := \ker(H^1(G_S, \mu_p) \longrightarrow \oplus_{v \in S} H^1(G_v, \mu_p)).$$

The failures of the isomorphism and duality in the tame case are reasons it is especially challenging.

DEFINITION 1.3. — *The set  $S$  of places  $K$  is called saturated if  $V_S/(\mathbb{K}^\times)^p = \{1\}$ .*

As consequence of Theorem 1.1, one has (see [13, Theorem 1.12])

THEOREM 1.4. — *A finite tame set  $S$  is saturated if and only if the Frobenius automorphisms  $\sigma_{\mathfrak{p}}$ ,  $\mathfrak{p} \in S$ , span the elementary  $p$ -abelian group  $\text{Gal}(K'(\sqrt[p]{V_\emptyset})/K')$ .*

We recall below the formula of Shafarevich applied in the case where  $S$  is tame (see for example [24, Chapter X, Section 7, Corollary 10.7.7]):

$$(1.1) \quad d(G_S) = |S| - d(\mathcal{O}_K^\times) + d(V_S/(\mathbb{K}^\times)^p).$$

Hence when  $S$  is saturated, one has  $d(G_S) = |S| - d(\mathcal{O}_K^\times)$ .

### 1.1.3. Spectral sequence

Let us start with the natural exact sequence

$$1 \longrightarrow H_S \longrightarrow G_S \longrightarrow G_\emptyset \longrightarrow 1,$$

where the group  $H_S$  is the closed normal subgroup of  $G_S$  generated by the tame inertia elements  $\tau_{\mathfrak{p}} \in G_S$ ,  $\mathfrak{p} \in S$ . Set

$$X_S := H_S/[H_S, G_\emptyset]H_S^p.$$

Recall as  $G_\emptyset$  is a pro- $p$  group, the compact ring  $\mathbb{F}_p[[G_\emptyset]]$  is local and acts continuously on  $H_S/[H_S, H_S]H_S^p$ . We give an easy lemma that can be found in [13] (see Lemmas 1.11 and 1.12).

LEMMA 1.5. — *Let  $S$  be a finite set of tame prime ideals of  $\mathcal{O}_K$ .*

- (i) *The  $\mathbb{F}_p[[G_\emptyset]]$ -module  $H_S/[H_S, H_S]H_S^p$  is topologically finitely generated by at most  $|S|$  elements.*
- (ii) *One has the exact sequence*

$$1 \longrightarrow H^1(G_\emptyset) \longrightarrow H^1(G_S) \longrightarrow X_S^\vee \longrightarrow \text{III}_\emptyset^2 \longrightarrow \text{III}_S^2.$$

*In particular, if  $S$  is such that  $H^1(G_\emptyset) \simeq H^1(G_S)$ , then  $X_S^\vee \hookrightarrow \text{III}_\emptyset^2$ . If moreover  $S$  is also saturated then  $X_S^\vee \simeq \text{III}_\emptyset^2$ .*

To conclude this subsection, let us observe the following: Let  $F_0/K_\emptyset$  be a cyclic extension of degree  $p$  in  $K_S/K$  such that  $F_0/K$  is Galois. Then  $F_0$  comes from a finite level: there exists a finite extension  $F/K$  and a cyclic extension  $F_1/F$  of degree  $p$ , ramified at some places above  $S$ , such that  $F_0 = K_\emptyset F_1$ . The estimate for  $\dim H^2(G_\emptyset)$  can be done by using the previous lemma, typically by seeking the fields  $F_0$ : this is the spirit of the method involving the Hochschild–Serre spectral sequence.

*Example 1.6 (The field  $\mathbb{Q}(\sqrt{-5460})$ ).* — Set  $p = 2$  and  $K = \mathbb{Q}(\sqrt{-5460})$ . The rational primes in  $\{43, 53, 101, 149, 157\}$  all split in  $K$ . Let  $S = \{\mathfrak{p}_{43}, \mathfrak{p}_{53}, \mathfrak{p}_{101}, \mathfrak{p}_{149}, \mathfrak{p}_{157}\}$ , the first primes above each of these as MAGMA computes them. We denote the abelian group  $\prod_{i=1}^d \mathbb{Z}/a_i$  by  $(a_1, \dots, a_d)$ . Computations show that:

- (i)  $\text{RCG}_K(\emptyset) = (2, 2, 2, 2)$ ;
- (ii)  $\text{RCG}_K(\mathfrak{p}_{53}, \mathfrak{p}_{101}, \mathfrak{p}_{149}, \mathfrak{p}_{157}) = (4, 8, 8, 8)$ ; Furthermore, for each of these primes  $\mathfrak{p}$  we compute  $\text{RCG}_K(\mathfrak{p}) = (2, 2, 2, 4)$ .
- (iii)  $\text{RCG}_K(S) = (8, 8, 8, 8)$ .

Since  $\mathcal{O}_K^\times = \pm 1$ , one easily sees  $d(V_\emptyset / (K^\times)^2) = 4 + 1 = 5$  hence  $S$  is saturated by (iii) and equality (1.1), and (ii) implies  $d(X_{\{\mathfrak{p}_{53}, \mathfrak{p}_{101}, \mathfrak{p}_{149}, \mathfrak{p}_{157}\}}) = 4$  and then  $d(X_S) \geq 4$ . As  $X_{\{\mathfrak{p}\}}$  is nontrivial for  $\mathfrak{p} \in \{\mathfrak{p}_{53}, \mathfrak{p}_{101}, \mathfrak{p}_{149}, \mathfrak{p}_{157}\}$ , we see there is a quadratic extension above  $K_\emptyset$  ramified at  $\mathfrak{p}$ . We have produced four independent elements of  $\text{III}_\emptyset^2$ .

Now take  $F = K(i) \subset K_\emptyset$ ;  $\mathfrak{p}_{43}$  is inert in  $F/K$ . An easy computation shows that  $\text{RCG}_F(\emptyset) = (4, 2, 2, 2)$  and  $\text{RCG}_F(\mathfrak{p}_{43}) = (4, 4, 2, 2)$ . As  $F_\emptyset = K_\emptyset$  we have an extension over  $K_\emptyset$  ramified only at  $\mathfrak{p}_{43}$ , so  $d(X_S) = 5$ , and by Lemma 1.5 we conclude that  $r(G_\emptyset) = 5$ .

### 1.2. Universal norms and relations

Put  $\mathcal{O}_{K_\emptyset}^\times = \bigcup_F \mathcal{O}_F^\times$ , where  $F/K$  run through the finite Galois extensions in  $K_\emptyset/K$ . Recall the following theorem due to Wingberg [40]; see also [24, Theorem 8.8.1, Chapter VIII, Section 8] where we take  $S = T = \emptyset$ ,  $\mathfrak{c}$  to be the full class of finite  $p$ -groups and  $A = \mathbb{Z}$ . We have written the results there in our notation.

**THEOREM 1.7 (Wingberg).** — *One has  $\widehat{H}^i(G_\emptyset, \mathcal{O}_{K_\emptyset}^\times) \simeq \widehat{H}^{3-i}(G_\emptyset, \mathbb{Z})^\vee$ .*

The exact sequence  $0 \rightarrow \mathbb{Z} \xrightarrow{\times p} \mathbb{Z} \rightarrow \mathbb{Z}/p \rightarrow 0$  gives:

$$0 \longrightarrow H^2(G_\emptyset, \mathbb{Z})/p \longrightarrow H^2(G_\emptyset, \mathbb{Z}/p) \longrightarrow H^3(G_\emptyset, \mathbb{Z})[p] \longrightarrow 0.$$

Taking the Pontryagin dual, one obtains:

$$(1.2) \quad 0 \longrightarrow H^3(G_\emptyset, \mathbb{Z})^\vee/p \longrightarrow H^2(G_\emptyset, \mathbb{Z}/p)^\vee \longrightarrow H^2(G_\emptyset, \mathbb{Z})^\vee[p] \longrightarrow 0.$$

By Theorem 1.7:

$$H^2(G_\emptyset, \mathbb{Z})^\vee \simeq H^1(G_\emptyset, \mathcal{O}_{K_\emptyset}^\times), \text{ and } H^3(G_\emptyset, \mathbb{Z})^\vee \simeq \widehat{H}^0(G_\emptyset, \mathcal{O}_{K_\emptyset}^\times).$$

Recall

$$\widehat{H}^0(G_\emptyset, \mathcal{O}_{K_\emptyset}^\times) \simeq \varprojlim_F \mathcal{O}_K^\times / N_{F/K} \mathcal{O}_F^\times,$$

where  $F/K$  run through the finite Galois extensions in  $K_\emptyset/K$ ,  $N_{F/K}$  is the norm in  $F/K$ , and  $H^1(G_\emptyset, \mathcal{O}_{K_\emptyset}^\times)$  is the  $p$ -part of  $\text{Cl}_K$  (see for example [24, Lemma 8.8.4, Chapter VIII, Section 8]).

This observation associated to Theorem 1.7 allows us to prove:

**COROLLARY 1.8.** — *One has  $\text{Def}(G_\emptyset) = d(\mathcal{E}_K/N_{K_\emptyset/K}\mathcal{E}_{K_\emptyset})$ , where  $N_{K_\emptyset/K}\mathcal{E}_{K_\emptyset} := \bigcap_{F/K} N_{F/K}\mathcal{E}_F$ . In particular when  $[F : K]$  is sufficiently large one has  $\text{Def}(G_\emptyset) = d(\mathcal{E}_K/N_{F/K}\mathcal{E}_F)$ .*

*Proof.* — If  $F/K$  is a finite Galois extension in  $K_\emptyset/K$  then  $(\mathcal{O}_K^\times)^{[F:K]} \subset N_{F/K}\mathcal{O}_F^\times$ , hence  $\mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times$  is a finite abelian  $p$ -group and  $\varprojlim_F \mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times$  is an abelian pro- $p$  group (obviously finitely generated). Then  $\varprojlim_F \mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times \simeq \varprojlim_F \mathbb{Z}_p \otimes (\mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times)$ . But as  $\mathbb{Z}_p$  is  $\mathbb{Z}$ -flat, one gets  $\mathbb{Z}_p \otimes (\mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times) \simeq E_K/(\mathbb{Z}_p \otimes N_{F/K}\mathcal{O}_F^\times) = E_K/\overline{N_{F/K}\mathcal{O}_F^\times}$ , where  $E_K = \mathbb{Z}_p \otimes \mathcal{O}_K^\times$  and  $\overline{N_{F/K}\mathcal{O}_F^\times}$  is the closure of  $N_{F/K}\mathcal{O}_F^\times$  in  $\mathbb{Z}_p \otimes \mathcal{O}_K^\times$ . Hence,

$$\varprojlim_F \mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times \simeq E_K / \bigcap_F \overline{N_{F/K}\mathcal{O}_F^\times}.$$

Thus

$$\mathbb{F}_p \otimes \varprojlim_F \mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times \simeq E_K/E_K^p \bigcap_F \overline{N_{F/K}\mathcal{O}_F^\times} \simeq \mathcal{E}_K/N_{K_\emptyset/K}\mathcal{E}_{K_\emptyset}.$$

The exact sequence (1.2) becomes

$$(1.3) \quad 0 \longrightarrow \mathcal{E}_K/N_{K_\emptyset/K}\mathcal{E}_{K_\emptyset} \longrightarrow H^2(G_\emptyset, \mathbb{Z}/p)^\vee \longrightarrow \text{Cl}_K[p] \longrightarrow 0,$$

and computing dimensions gives the result.  $\square$

For  $\#G_\emptyset < \infty$  it has been known for a long time that the number of relations of  $G_\emptyset$  is related to the norm of the units in the tower. See for example Section 2 of [28].

As a consequence, one also has

**COROLLARY 1.9.** — *Let  $F/K$  be a finite Galois extension in  $K_\emptyset/K$ . Then  $\text{Def}(G_\emptyset) \geq d(\mathcal{O}_K^\times/N_{F/K}\mathcal{O}_F^\times)$ , and one has equality when  $F$  is sufficiently large.*

*Proof.* — This is obvious using that  $\mathcal{E}_K/N_{K_\emptyset/K}\mathcal{E}_{K_\emptyset} \twoheadrightarrow \mathcal{E}_K/N_{F/K}\mathcal{E}_F$ . For the equality, use the fact that  $\mathcal{E}_K$  is finite.  $\square$

When  $p = 2$ , if  $-1$  is not a norm of a unit in a quadratic subextension  $F/K$  of  $K_\emptyset/K$ , then  $-1 \notin N_{K_\emptyset/K}\mathcal{O}_{K_\emptyset}^\times$ , which implies  $\text{Def}(G_\emptyset) \geq 1$ . We will see that this condition appears almost all the time when  $K$  is an imaginary quadratic extension. We close this subsection with a basic fact.

**FACT.** — *For  $S$  a finite set of tame places,  $0 \leq \text{Def}(G_S)$ .*

*Proof.* — We refer to [27], especially Lemma 6.8.6, for the facts we need concerning the homology of profinite groups. From the exact sequence of compact groups

$$0 \longrightarrow \mathbb{Z}_p \xrightarrow{\times p} \mathbb{Z}_p \longrightarrow \mathbb{Z}/p \longrightarrow 0$$

we obtain the homology sequence

$$\cdots \longrightarrow H_2(G_S, \mathbb{Z}/p) \longrightarrow H_1(G_S, \mathbb{Z}_p)[p].$$

As  $H_1(G_S, \mathbb{Z}_p) \simeq G_S^{ab}$ , we have  $d(G_S) = d(G_S^{ab}[p]) \leq d(H_2(G_S, \mathbb{Z}/p)) = r(G_S)$ .  $\square$

### 1.3. Minkowski units

#### 1.3.1.

Recall that for a finite group  $G$ , the ring  $\mathbb{F}_p[G]$  is a Frobenius algebra (see for example [5, Section 62]): every free submodule of an  $\mathbb{F}_p[G]$ -module  $M$  is in direct sum so we may write  $M = \mathbb{F}_p[G]^t \oplus N$ , where  $N$  is torsion (for every element  $n \in N$ , there exists  $0 \neq h \in \mathbb{F}_p[G]$  such that  $h \cdot n = 0$ ), and  $t := t_G(M)$  is uniquely determined (by Krull-Schmidt Theorem). Observe that if  $M^\wedge$  is the Pontryagin dual of  $M$ , then  $t_G(M) = t_G(M^\wedge)$ .

We record some useful properties. Let  $H \subset G$  be a subgroup of  $G$ .

- (i) Recall first that by Mackey's decomposition theorem, one has the isomorphism of  $\mathbb{F}_p[H]$ -modules  $\text{Res}_H \mathbb{F}_p[G] \simeq \mathbb{F}_p[H]^{[G:H]}$ .
- (ii) Suppose moreover  $H \triangleleft G$ , and denote by  $N_H = \sum_{h \in H} h \in \mathbb{F}_p[G]$  the norm map from  $H$ . For an  $\mathbb{F}_p[G]$ -module  $M$  let  $M^H$  denote the invariants. Then one easily obtains the isomorphism of  $\mathbb{F}_p[G]$ -modules

$$(1.4) \quad \mathbb{F}_p[G/H] \simeq \mathbb{F}_p \otimes_{\mathbb{F}_p[H]} \mathbb{F}_p[G] \simeq \mathbb{F}_p[G]^H$$

and  $N_H(\mathbb{F}_p[G]) = \mathbb{F}_p[G]^H$  so

$$(1.5) \quad N_H(\mathbb{F}_p[G]) \simeq \mathbb{F}_p[G/H]$$

as  $\mathbb{F}_p[G/H]$ -modules.

#### 1.3.2.

Let  $F/K$  be a finite Galois extension of number fields with Galois group  $G$ .

DEFINITION 1.10. — Let  $\mathcal{E}_F := \mathbb{F}_p \otimes \mathcal{O}_F^\times$ . We say that  $F/K$  has a Minkowski unit (at  $p$ ), if  $\mathcal{E}_F$  contains a nontrivial free  $\mathbb{F}_p[G]$ -submodule. In other word,  $F/K$  has a Minkowski unit if  $t_G(\mathcal{E}_F) \geq 1$ .

Hence the quantity  $t_G(\mathcal{E}_F)$  measures “the number” of independent Minkowski units in  $F/K$ .

If  $(p, |G|) = 1$  then  $\mathcal{E}_F$  is a semisimple  $\mathbb{F}_p[G]$ -module. Determining the existence of Minkowski units is more difficult when  $(p, |G|) = p$ . When  $G$  is a  $p$ -group, and  $F/K$  is unramified, it is tempting to regard the existence of a Minkowski unit in  $F/K$  as rare.

### 1.3.3. Example

We want to illustrate the notion of Minkowski units.

LEMMA 1.11. — Let  $F/K$  be a  $p$ -extension with Galois group  $G$ . Let  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_k\}$  be a set of tame primes of  $K$  that split completely in  $F/K$ . If  $d(G_{F,S}) = d(G_{F,\emptyset})$  then  $t_G(V_{F,\emptyset}) \geq k$ , and if  $|G_{F,S}^{ab}| = |G_{F,\emptyset}^{ab}|$  then  $t_G(\mathcal{E}_F) \geq k$ .

*Proof.* — Observe first that  $G$  acts on  $\text{Gal}(F'(\sqrt[p]{V_{F,\emptyset}})/F')$  (resp. on  $\text{Gal}(F'(\sqrt[p]{\mathcal{O}_F^\times})/F')$ ). Then by Remark 1.2, one has

$$(1.6) \quad t_G(V_F/(F^\times)^p) = t_G(\text{Gal}(F'(\sqrt[p]{V_{F,\emptyset}})/F')),$$

and

$$(1.7) \quad t_G(\mathcal{E}_F) = t_G((\mathcal{E}_F)^\wedge) = t_G(\text{Gal}(F'(\sqrt[p]{\mathcal{O}_F^\times})/F')).$$

For all prime ideals  $\mathfrak{P}_{ij}|\mathfrak{p}_i$  of  $\mathcal{O}_F$  we consider the Frobenius automorphisms  $\sigma_{\mathfrak{P}_{ij}}$  in  $\text{Gal}(F'(\sqrt[p]{V_{F,\emptyset}})/F')$ . Note that we are no longer in an abelian over  $K$  situation as just before Theorem 1.1. By Theorem 1.1 (i), the hypothesis  $d(G_{F,S}) = d(G_{F,\emptyset})$  implies the Frobenius automorphisms  $\sigma_{\mathfrak{P}_{ij}}$  in  $\text{Gal}(F'(\sqrt[p]{V_{F,\emptyset}})/F')$  are independent. As each  $\mathfrak{p}_i$  splits completely in  $F/K$ , we have that  $\text{Gal}(F'(\sqrt[p]{V_{F,\emptyset}})/F')$  contains  $k$  distinct free  $\mathbb{F}_p[G]$ -modules, one for each  $\mathfrak{p}_i$ . The first assertion follows by (1.6). For the second assertion, use the second part of Theorem 1.1 and (1.7). □

Recall that we denote the abelian group  $\prod_{i=1}^d \mathbb{Z}/a_i\mathbb{Z}$  by  $(a_1, \dots, a_d)$ . Let  $p = 2$  and  $K = \mathbb{Q}(\sqrt{5 \cdot 13 \cdot 17 \cdot 29})$  and let  $H = \mathbb{Q}(\sqrt{5}, \sqrt{13}, \sqrt{17}, \sqrt{29})$  be its Hilbert class field. Here  $\text{Cl}_K = (2, 2, 2)$ , and  $\text{Cl}_H = (4, 4)$ . Consider the primes  $\ell = 2311$  and  $q = 3319$ . We easily see  $\ell\mathcal{O}_K = \mathfrak{l}_1\mathfrak{l}_2$  and  $q\mathcal{O}_K = \mathfrak{q}_1\mathfrak{q}_2$  and these ideals are all principal. In the table below we compute the 2-parts of the ray class groups for  $K$  and  $H$  of the given conductors. The

Table 1.1. Ray Class Groups

Conductor	K	H
1	(2, 2, 2)	(4, 4)
$\mathfrak{l}_i, i \in \{1, 2\}$	(2, 2, 2)	(4, 4)
$\mathfrak{q}_i, i \in \{1, 2\}$	(2, 2, 2)	(4, 4)
$\mathfrak{l}_1\mathfrak{q}_1$	(2, 2, 2)	(2, 2, 2, 4, 4)
$\mathfrak{l}_1\mathfrak{q}_2$	(2, 2, 2, 2)	(2, 2, 2, 2, 2, 4, 8)
$\mathfrak{l}_2\mathfrak{q}_1$	(2, 2, 2, 2)	(2, 2, 2, 2, 2, 4, 8)
$\mathfrak{l}_2\mathfrak{q}_2$	(2, 2, 2)	(2, 2, 2, 4, 4)

computations were done with MAGMA (see [3]) and assume the GRH.

Note that in the first three rows, the ray class groups are identical. As the principal ideals  $\mathfrak{l}_i$  and  $\mathfrak{q}_i$  split completely in  $H/K$ , by Lemma 1.11 one sees that  $\mathcal{O}_H^\times \otimes \mathbb{F}_2$  has a Minkowski unit over  $K$ : in other words putting  $G = \text{Gal}(H/K) \simeq (\mathbb{Z}/2\mathbb{Z})^3$ , one has  $t_G(\mathcal{E}_H) \geq 1$ . Note  $H$  is a degree 16 totally real field so  $\dim \mathcal{E}_H = 16$  and  $\mathcal{E}_H \simeq_G \mathbb{F}_2[G] \oplus M$  where  $\dim M = 8$  so  $M$  a priori could be free.

We now show that  $M$  is not free. Set  $K_0 = K$ ,  $K_1 = \mathbb{Q}(\sqrt{5 \cdot 17})$ , and  $K_2 = \mathbb{Q}(\sqrt{13 \cdot 29})$ . Let  $F$  be the biquadratic field  $K_1K_2$ . Computations show that  $\text{Cl}_{K_1} = (2)$ ,  $\text{Cl}_{K_2} = (2)$ , and  $\text{Cl}_F = (2, 4)$ . Denote by  $\varepsilon_i$  the fundamental unit of  $K_i$ , and put

$$e = \# (\mathcal{O}_F^\times / \langle -1, \varepsilon_i, i = 0, 1, 2 \rangle).$$

Applying the Brauer class number formula in the biquadratic extension  $F/\mathbb{Q}$ , i.e.  $|\text{Cl}_F| = \frac{1}{4}e|\text{Cl}_K||\text{Cl}_{K_1}||\text{Cl}_{K_2}|$ , to deduce  $e = 1$ , and then  $\mathcal{O}_F^\times = \langle -1, \varepsilon_i, i = 0, 1, 2 \rangle$ .

Let  $\sigma$  be a generator of  $G = \text{Gal}(F/K)$ . We compute:

- (i) the norm of  $\varepsilon_2$  in  $F/K$  is  $+1$ ,
- (ii) the norm of  $\varepsilon_1$  in  $F/K$  is  $-1$ ,
- (iii)  $\sigma$  acts trivially on  $\varepsilon_0$ ,

Hence, as we will observe in Lemma 5.1, one obtains that  $\mathcal{E}_F \simeq \mathbb{F}_2[G'] \oplus \mathbb{F}_2^2$ , where  $G' = \text{Gal}(F/K)$ . Finally since  $\mathcal{E}_H \twoheadrightarrow \mathcal{E}_F$ , and  $t_{G'}(\mathcal{E}_F) \geq t_G(\mathcal{E}_H)$  we conclude that  $t_G(\mathcal{E}_H) = 1$ .

## 2. Detecting the relations along $K_\emptyset/K$

As mentioned in the remark in the introduction, we can easily find  $d$  elements of  $\text{III}_\emptyset^2$  by constructing ramified extensions at a low level in the



tower  $K_\theta/K$ . For  $(G_n)$  a sequence of open normal subgroups of  $G$  with  $\bigcap_{n=1}^\infty G_n = \{e\}$ , let  $K_n$  be the fixed field of  $G_n$  and set  $H_n = \text{Gal}(K_n/K)$ . In this section we show that the presence of torsion elements in the  $\mathbb{F}_p[H_n]$ -module  $\text{Gal}(K'_n(\sqrt[p]{\mathcal{O}_{K_n}^\times})/K'_n)$  can give rise to more relations.

### 2.1. First observations

Let  $p$  be a prime number and  $K$  be a number field.

If  $F/K$  is a Galois extension with Galois group  $G$ , the norm map  $N_G$  sends  $\mathcal{E}_F$  to  $\frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p} \subset \mathcal{E}_F$ ; denote by  $N'_G : \mathcal{E}_F \rightarrow \mathcal{E}_K$  the map from  $\mathcal{E}_F$  to  $\mathcal{E}_K$  induced by the norm in  $F/K$ . The commutative diagram:

$$\begin{array}{ccc}
 \mathcal{E}_F & \xrightarrow{N_G} & \frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p} \hookrightarrow \mathcal{E}_F \\
 & \searrow N'_G & \uparrow \\
 & & \mathcal{E}_K
 \end{array}$$

implies the following easy lemma:

LEMMA 2.1. — *One has  $N'_G(\mathcal{E}_F) \twoheadrightarrow N_G(\mathcal{E}_F)$ . Moreover,  $\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p = (\mathcal{O}_K^\times)^p \implies N'_G(\mathcal{E}_F) \simeq N_G(\mathcal{E}_F)$ .*

The study of the norm map  $N_G$  is “purely algebraic”, i.e. it does not involve number theory. Lemma 2.2 below is proved at the beginning of the proof of [25, Lemma 2]). Since that Lemma is stated differently we include a proof that is essentially from [25].

LEMMA 2.2. — *Let  $G$  be a finite  $p$ -group and  $M$  an  $\mathbb{F}_p[G]$ -module. Let  $N_G : M \rightarrow M$  be the norm map. Let  $m \in M$ . Then  $N_G(m) = 0$  if and only if  $m$  is a torsion element.*

*Proof.* — Let  $0 \neq m \in M$ . Recall that the annihilator  $A_m$  of a nontrivial element  $m \in M$  is an ideal of  $\mathbb{F}_p[G]$  and that  $m$  is a torsion element if and only if  $A_m \neq 0$ .

If  $A_m = 0$  then the  $\mathbb{F}_p[G]$ -span of  $m$  is isomorphic to  $\mathbb{F}_p[G]$  and as  $N_G(\mathbb{F}_p[G]) = \mathbb{F}_p$  by (1.5), we see  $N_G(m) \neq 0$ .

Conversely, suppose that  $A_m \neq 0$ . Then  $A_m^G \neq 0$  since  $G$  is a  $p$ -group acting on a nontrivial  $\mathbb{F}_p$ -vector space. Hence  $A_m^G \subset (\mathbb{F}_p[G])^G$  which is in turn the one-dimensional vector space  $\mathbb{F}_p \cdot N_G$ . Thus  $\mathbb{F}_p \cdot N_G = A_m^G \subset A_m$  so  $N_G(m) = 0$ . □

More generally, one has

**THEOREM 2.3.** — *Let  $F/K$  be a finite  $p$ -extension with Galois group  $G$  and write  $N'_G(\mathcal{E}_F) \simeq \mathbb{F}_p^t$ . Then  $t_G(\mathcal{E}_F) \leq t \leq t_G(\mathcal{E}_F) + d\left(\frac{\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p}{(\mathcal{O}_K^\times)^p}\right)$ . In particular if  $\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p = (\mathcal{O}_K^\times)^p$ , then  $t = t_G(\mathcal{E}_F)$ .*

*Proof.* — Write  $\mathcal{E}_F \simeq \mathbb{F}_p[G]^{t_G(\mathcal{E}_F)} \oplus N$ , where  $N$  is generated by torsion elements as an  $\mathbb{F}_p[G]$ -module. By (1.5) and Lemma 2.2 one has  $N_G(\mathcal{E}_F) \simeq \mathbb{F}_p^{t_G(\mathcal{E}_F)}$ . So by Lemma 2.1 we see  $N'_G(\mathcal{E}_F) \simeq N_G(\mathcal{E}_F) \simeq \mathbb{F}_p^{t_G(\mathcal{E}_F)}$ , proving the result when  $\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p = (\mathcal{O}_K^\times)^p$ .

By noting that the ‘difference’ between  $N_G(\mathcal{E}_F)$  and  $N'_G(\mathcal{E}_F)$  is exactly  $\frac{N_G(\mathcal{O}_F^\times) \cap (\mathcal{O}_F^\times)^p}{(\mathcal{O}_K^\times)^p}$  which has  $p$ -rank at most  $d\left(\frac{\mathcal{O}_K^\times \cap (\mathcal{O}_F^\times)^p}{(\mathcal{O}_K^\times)^p}\right)$ , we obtain the general case. □

### 2.2. Exhibiting relations via the Hochschild–Serre spectral sequence

In this subsection, we flesh out the details of the process described in the Remark of the Introduction for explicitly exhibiting  $d(G_\emptyset)$  relations in a minimal presentation of  $G_\emptyset$ . It would be helpful to refer to Figure 0.1 from the introduction. Put  $K' = K(\zeta_p)$ . Let  $S = S_1 \cup S_2$  be a set of tame prime ideals of  $\mathcal{O}_K$  such that:

- (1)  $S_1$  is a minimal set whose Frobenius automorphisms generate the  $d(G_\emptyset)$ -dimensional  $\mathbb{F}_p$ -vector space  $\text{Gal}(K'(\sqrt[p]{\mathcal{V}_\emptyset})/K'(\sqrt[p]{\mathcal{O}_K^\times}))$ , and
- (2)  $S_2$  is a minimal set whose Frobenius automorphisms generate the  $\mathbb{F}_p$ -vector space  $\text{Gal}(K'(\sqrt[p]{\mathcal{O}_K^\times})/K')$  of dimension  $r_1 + r_2 - 1 + \delta$ .

Recall the Frobenius automorphisms above are well-defined up to nonzero scalar multiples in the Galois groups, which are vector spaces over  $\mathbb{F}_p$ . This ambiguity does not affect their spanning properties. One has

**LEMMA 2.4.** — *The set  $S$  is saturated, in particular  $\text{III}_S^2 = \{0\}$ . Moreover  $d(G_S) = d(G_\emptyset)$ , and  $r(G_\emptyset) = d(X_S)$ .*

*Proof.* — That  $S$  is saturated follows immediately from Theorem 1.4. As there is no dependence relation between the Frobenius automorphisms (the set  $S$  is minimal), Theorem 1.1 implies  $d := d(G_S) = d(G_\emptyset)$ . That  $r(G_\emptyset) = d(X_S)$  follows from the second part with Lemma 1.5. □

**LEMMA 2.5.** — *Write  $(a_1, \dots, a_d)$  for the  $p$ -part of  $\text{RCG}_K(\emptyset)$  and let  $S_1 = \{\mathfrak{p}_1, \dots, \mathfrak{p}_d\}$  as above. Then  $\text{RCG}_K(\mathfrak{p}_1, \dots, \mathfrak{p}_d) \twoheadrightarrow (pa_1, \dots, pa_d)$ .*

*Proof.* — This is a consequence of Theorem 1.1. As the primes of  $S_1$  split completely in the governing extension  $\text{Gal}(K'(\sqrt[p]{\mathcal{O}_K^\times})/K')$ , for each prime ideal  $\mathfrak{p} \in S_1$  we have  $\#G_{\{\mathfrak{p}\}}^{ab} \neq \#G_\emptyset^{ab}$ . We conclude by noting that  $d(G_{S_1}) = d(G_\emptyset)$ .  $\square$

Lemma 2.5 implies the existence of  $d$  independent degree- $p$  cyclic extensions  $F_i$  of  $K_\emptyset$ , each totally ramified at  $\mathfrak{p}_i$ ,  $i = 1, \dots, d$ , and on which  $G_\emptyset$  acts trivially, implying that  $d(X_S) \geq d$ . *The rest of the relations are difficult and detected via the set  $S_2$ .*

### 2.3. Proof of Theorem A

Let  $(G_n)$  be a sequence of open normal subgroups of  $G_\emptyset$  such that  $G_n \subset G_{n+1}$  and  $\bigcap_n G_n = \{e\}$ . Put  $H_n := G_\emptyset/G_n$ ,  $K_n := K_\emptyset^{G_n}$ , and write  $\mathcal{E}_{K_n} := \mathbb{F}_p[\mathbf{H}_n]^{t_n} \oplus N_n$  where  $N_n$  is torsion as an  $\mathbb{F}_p[\mathbf{H}_n]$ -module.

LEMMA 2.6. — *The sequence  $(t_n)$  is nonincreasing.*

*Proof.* — Recall from (1.4) that the norm map from  $H_{n+1,n} := \text{Gal}(K_{n+1}/K_n)$  on  $\mathbb{F}_p[\mathbf{H}_{n+1}]$  induces the following  $\mathbb{F}_p[\mathbf{H}_n]$ -isomorphisms:

$$\mathbb{F}_p[\mathbf{H}_n] \simeq \mathbb{F}_p[\mathbf{H}_{n+1}]_{H_{n+1,n}} \simeq \mathbb{F}_p[\mathbf{H}_{n+1}]^{H_{n+1,n}}.$$

The norm map  $N_{H_{n+1,n}}$  of  $K_{n+1}/K_n$  induces a morphism from  $\mathcal{E}_{K_{n+1}}$  to  $\mathcal{E}_{K_n}$  which allows us to obtain

$$\mathbb{F}_p[\mathbf{H}_n]^{t_{n+1}} \hookrightarrow \frac{\mathcal{O}_{K_n}^\times}{\mathcal{O}_{K_n}^\times \cap (\mathcal{O}_{K_{n+1}}^\times)^p} \leftarrow \mathcal{E}_{K_n},$$

which implies  $t_n \geq t_{n+1}$ .  $\square$

DEFINITION 2.7. — *Set  $\lambda := \lambda_{K_\emptyset/K} = \lim_n t_n$ . We call this the Minkowski-rank of the units along  $K_\emptyset/K$ .*

One easily sees that  $\lambda$  does not depend on the sequence  $(G_n)$ .

Let us write  $p^\beta := [K'(\sqrt[p]{\mathcal{O}_K^\times}) \cap K'K_\emptyset : K'] = [\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\emptyset}^\times)^p : (\mathcal{O}_K^\times)^p]$ . Obviously,  $\beta \leq \min(d(\mathcal{O}_K^\times), d(G_\emptyset))$ .

PROPOSITION 2.8. — *One has:  $\delta = 0 \implies \beta = 0$ .*

*Proof.* — Recall  $K' := K(\zeta_p)$ . Let  $\Delta = \text{Gal}(K'/K)$  be the Galois group of  $K'/K$ . As  $\Delta$  acts trivially on  $\mathcal{O}_K^\times$ , by Kummer duality the action of  $\Delta$  over  $\text{Gal}(K'(\sqrt[p]{\mathcal{O}_K^\times})/K')$  is given by the cyclotomic character; in particular, there is no nontrivial subspace of  $\text{Gal}(K'(\sqrt[p]{\mathcal{O}_K^\times})/K')$  on which  $\Delta$  acts trivially. As  $\Delta$  acts trivially on  $\text{Gal}(K'K_\emptyset/K')$ , the result holds.  $\square$

THEOREM 2.9. — We have the estimates:

$$d(\mathcal{O}_K^\times) - \lambda - \beta \leq \text{Def}(G_\emptyset) \leq d(\mathcal{O}_K^\times) - \lambda.$$

In particular,

- if  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\emptyset}^\times)^p = (\mathcal{O}_K^\times)^p$  or if  $\delta = 0$ , then  $\text{Def}(G_\emptyset) = d(\mathcal{O}_K^\times) - \lambda$ .
- if  $\lambda = d(\mathcal{O}_K^\times)$  then  $\text{Def}(G_\emptyset) = 0$ .

*Proof.* — We keep the notations of the beginning of the section.

We give two proofs for the lower bound. The first one is ‘algebraic’ while the second is number-theoretic and is more ‘explicit’ in how we determine the existence of the relations.

We first establish the upper bound. Denote by  $N_{H_n}$  the norm map for the extension  $K_n/K$ . Observe that by Corollary 1.8,  $\text{Def}(G_\emptyset) = d(\mathcal{E}_K) - d(N'_{H_n}(\mathcal{E}_{K_n}))$  for  $n \gg 0$ . Take  $n$  sufficiently large such that  $t_n = \lambda$ . One has  $d(N_{H_n}(\mathcal{E}_{K_n})) \geq \lambda$  (see Theorem 2.3), implying that  $d(N'_{H_n}(\mathcal{E}_{K_n})) \geq \lambda$ . Hence one gets:

$$\text{Def}(G_\emptyset) \leq d(\mathcal{O}_K^\times) - \lambda.$$

Below are the two proofs of the lower bound.

*First proof.* — Observe that  $\beta = d\left(\frac{\mathcal{O}_K^\times \cap (\mathcal{O}_{K_n}^\times)^p}{(\mathcal{O}_K^\times)^p}\right)$  since  $n \gg 0$ . By Theorem 2.3 one also has  $d(N'_{H_n}(\mathcal{E}_{K_n})) \leq \lambda + \beta$ , and then by Corollary 1.8 we get

$$\text{Def}(G_\emptyset) \geq d(\mathcal{O}_K^\times) - \lambda - \beta.$$

*Second proof.* — Here we show  $d(\mathcal{O}_K^\times) - \lambda - \beta \leq \text{Def}(G_\emptyset)$  using saturated sets and the Hochschild–Serre exact sequence.

First assume that  $\zeta_p \in K$  i.e.  $\delta = 1$ . Choose  $n \gg 0$ , and write  $\mathcal{E}_{K_n} = \mathbb{F}_p[\mathbb{H}_n]^\lambda \oplus N_n$ , where  $N_n$  is a  $H_n$ -torsion  $\mathbb{F}_p[\mathbb{H}_n]$ -module.

$$\text{Put } \mathcal{E}'_K := \frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap (\mathcal{O}_{K_n}^\times)^p} \hookrightarrow \mathcal{E}_{K_n}.$$

Hence  $[L_4 : K_n] = \#\mathcal{E}'_K$ . Observe that  $\text{Gal}(L_4/K_n) \simeq \mathbb{F}_p^t$ , where  $t = d(\mathcal{O}_K^\times) - \beta$ .

We will find a set of primes of  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_{t-\lambda}\}$  in  $K$  such that:

- $\mathfrak{p}_i$  splits completely in  $K_n/K$ ,
- Their Frobenius automorphisms span a  $(t - \lambda)$ -dimensional space in  $\text{Gal}(L_2/L_1) \simeq \text{Gal}(L_4/K_n)$ ,
- For each  $i$ , let  $\mathfrak{b}_{ij}$  be the primes above  $\mathfrak{p}_i$  in  $K_n$ . There is a dependence relation on the Frobenius automorphisms of the  $\mathfrak{b}_{ij}$  in  $\text{Gal}(L_5/K_n)$ . By Gras–Munnier (Theorem 1.1) this implies the existence of a  $\mathbb{Z}/p$ -extension  $R_i/K_n$  ramified only at (these primes




Figure 2.1.

above)  $\mathfrak{p}_i$ . Let  $\tilde{R}_i$  be the Galois closure over  $K$  of  $R_i$ . As the  $p$ -group  $\text{Gal}(K_n/K)$  must act on the  $\mathbb{F}_p$ -vector space  $\text{Gal}(\tilde{R}_i/K_n)$  with a fixed point, by iteration we may assume  $R_i/K$  is Galois. The  $\mathbb{Z}/p$ -extension  $R_iK_\emptyset$  is ramified only at  $\mathfrak{p}_i$  and gives an element of  $\text{III}_\emptyset^2$ . We have produced  $t - \lambda$  elements of  $\text{III}_\emptyset^2$  in addition to the  $d$  elements of  $\text{III}_\emptyset^2$  we get by choosing primes  $\{\mathfrak{q}_1, \dots, \mathfrak{q}_d\}$  of  $K$  whose Frobenius automorphisms form a basis of  $\text{Gal}(L_3/L_2)$ .

This gives the lower bound. We now construct  $S$ .

As  $L_2/K$  is abelian ( $\zeta_p \in K$ ),  $H_n = \text{Gal}(K_n/K)$  acts trivially on  $\text{Gal}(L_2/L_1)$  (and thus on  $\text{Gal}(L_4/K_n)$  as well).

After taking the Kummer dual of  $\mathcal{E}_{K_n}$ , one obtains  $\text{Gal}(L_5/K_n) \simeq \mathbb{F}_p[H_n]^\lambda \oplus M_n$ , where  $M_n = N_n^\vee$  is a  $H_n$ -torsion  $\mathbb{F}_p[H_n]$ -module. The natural surjection  $\pi : \text{Gal}(L_5/K_n) \twoheadrightarrow \text{Gal}(L_4/K_n)$  induces, upon taking  $H_n$ -coinvariants, the map

$$\text{Gal}(L_5/K_n)_{H_n} \simeq \mathbb{F}_p^\lambda \oplus (M_n)_{H_n} \xrightarrow{\pi} \text{Gal}(L_4/K_n) \simeq \text{Gal}(L_2/L_1) \simeq \mathbb{F}_p^t.$$

Thus

$$d(\pi((M_n)_{H_n})) \geq t - \lambda = d(\mathcal{O}_K^\times) - \beta - \lambda.$$

Take (at least)  $t - \lambda$  elements  $x_i$  in  $\text{Gal}(L_5/K_n)$ , such that their image under the projection  $\pi$  forms a basis of  $\pi((M_n)_{H_n}) \subset \text{Gal}(L_4/K_n) \simeq \text{Gal}(L_2/L_1) \simeq \mathbb{F}_p^t$ . We choose  $\mathfrak{p}_i$  to split completely in  $K_n/K$  and have Frobenius  $\pi(x_i) \in \text{Gal}(L_2/L_1)$ , so clearly  $\mathfrak{p}_i$  satisfies the first two points above. We have chosen  $\mathfrak{p}_i$  so that the primes above it in  $K_n$  have Frobenius automorphisms generating a  $\mathbb{F}_p[H_n]$ -torsion module in  $\text{Gal}(L_5/K_n)$ . This settles the third point and the case  $\delta = 1$ .

Now suppose  $\delta = 0$ . Replace every field  $L_i$  above by  $L'_i := L_i(\zeta_p)$ . The key fact is this: by Proposition 2.8, one has  $d(\text{Gal}(L'_2/L'_1)) = d(\mathcal{O}_K^\times)$  so  $L'_2 \cap K_n = K$ . This disjointness allows us to apply the Chebotarev density theorem as above. The rest of the proof is word for word the same from this point on.

The last result follows since  $\text{Def}(G_\theta) \geq 0$ . □

*Remark 2.10.* — Observe that

- (i) the inequality  $\text{Def}(G_\theta) \leq d(\mathcal{O}_K^\times) - \lambda$  comes from universal norms of units and is Wingberg's result (Theorem 1.7);
- (ii) the group  $G_\theta$  has at least  $\lambda$  fewer relations than the maximal possible number,  $\dim V_\theta/(K^\times)^p$ .

**COROLLARY 2.11.** — *Suppose  $K_\theta/K$  is finite. Then  $\lambda < d(\mathcal{O}_K^\times) - d^2/4 + d$ .*

*Proof.* — By the Theorem of Golod–Shafarevich one has  $\text{Def}(G_\theta) > d^2/4 - d$ ; then apply Theorem 2.9. □

### 2.4. Remarks when $G_\theta$ is abelian

• Consider first the case where  $G_\theta$  is cyclic. Clearly  $d(G) = r(G) = 1$  so  $\text{Def}(G_\theta) = 0$ . By Theorem 2.3, we get

$$\lambda = t_{G_\theta}(\mathcal{E}_{K_\theta}) \geq d(\mathcal{O}_K^\times) - \beta \geq d(\mathcal{O}_K^\times) - 1,$$

due to the fact that  $\beta \leq 1$ . In particular, this situation forces  $K_\theta$  to have a Minkowski unit provided  $K$  is neither  $\mathbb{Q}$  nor imaginary quadratic. We can recover this fact by using the well-known following result: as  $G_\theta$  is cyclic, every element of  $\mathcal{O}_K^\times$  is the norm of an element of  $\mathcal{O}_{K_\theta}^\times$ . Note this last argument applies in the quadratic imaginary case as well.

As an example, take the imaginary quadratic number field  $K = \mathbb{Q}(\sqrt{-q \cdot \ell})$ , with  $-q \equiv \ell \equiv 1 \pmod{4}$ . Here,  $p = 2$ ,  $G_\theta$  is cyclic, and  $\mathcal{O}_K^\times \cap \mathcal{O}_{K_\theta}^{\times 2} = \mathcal{O}_K^{\times 2}$ . We find  $\lambda = 1$ , and finally that  $\mathcal{E}_{K_\theta} \simeq \mathbb{F}_2[G_\theta]$ .

Observe that if  $G_\emptyset \simeq \mathbb{Z}/2\mathbb{Z}$ , then the fundamental unit of the biquadratic extension  $K(\sqrt{\ell})$  is exactly the fundamental unit of the quadratic field  $\mathbb{Q}(\sqrt{\ell})$  and then is of norm  $-1$ . We again have  $\mathcal{E}_{K_\emptyset} \simeq \mathbb{F}_2[G_\emptyset]$ .

- Take  $p = 2$ , and  $K$  such that  $G_\emptyset \simeq (\mathbb{Z}/2\mathbb{Z})^2$ . Here  $d(G) = 2$  and  $r(G) = 3$  so  $\text{Def}(G_\emptyset) = 1$ , implying the existence of a difficult relation. By Theorem 2.3, we get

$$\lambda = t_{G_\emptyset}(\mathcal{E}_{K_\emptyset}) \geq d(\mathcal{O}_K^\times) - 2,$$

due to the fact that  $\beta \leq 2$ .

Let us be more precise: Kisilevsky in [14] showed that if  $G_\emptyset \simeq (\mathbb{Z}/2\mathbb{Z})^2$ , then for every quadratic subextension  $F_i/K$  in  $K_\emptyset/K$ , one has  $(\mathcal{O}_K^\times : N_{F_i/K} \mathcal{O}_{F_i}^\times) = 2$ . We prove

**PROPOSITION 2.12.** — *Let  $K/\mathbb{Q}$  be a quadratic extension such that  $G_\emptyset \simeq (\mathbb{Z}/2\mathbb{Z})^2$ . Then the difficult relation is detected at one of the three quadratic subextensions  $F/K$  in  $K_\emptyset/K$ .*

*Proof.* — Suppose first that  $K/\mathbb{Q}$  is an imaginary quadratic extension. By Kisilevsky’s result  $-1$  is not a norm of any unit in each of the three subextensions  $F_i/K$  of  $K_\emptyset/K$ . Let us choose  $F := F_i$  such that  $F \neq K(\sqrt{-1})$ ; put  $G = \text{Gal}(F/K)$ . By using  $N_G(\mathcal{O}_F^\times) \subset \{\pm 1\}$ , it is then easy to see that, modulo  $\mathcal{O}_F^{\times 2}$ ,  $-1$  is not a norm of any unit in  $F/K$  implying that the norm map  $N_G : \mathcal{E}_F \rightarrow \frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap \mathcal{O}_F^{\times 2}}$  is not onto.

Recall that  $\frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap \mathcal{O}_F^{\times 2}} \hookrightarrow \mathcal{E}_F$ . As  $\dim_{\mathbb{F}_2} \mathcal{E}_F = 2$ , the only possibilities for the  $\mathbb{F}_2[G]$ -module  $\mathcal{E}_F$  are  $\mathbb{F}_2^2$  and  $\mathbb{F}_2[G]$ . As the norm map is onto in the latter case we see  $\mathcal{E}_F \simeq \mathbb{F}_2^2$  and then  $t_G(\mathcal{E}_F) = 0$  so  $\lambda = 0$ : the difficult relation is detected by the quadratic extension  $F/K$ .

We now settle the case where  $K/\mathbb{Q}$  is real quadratic extension. Denote by  $\varepsilon$  the positive fundamental unit of  $K$ . By Kisilevsky’s result, one knows that for every quadratic subextension  $F_i/K$ ,  $-1$  or  $\varepsilon$  is not a norm of any unit in  $F_i/K$ . Take one such quadratic extension  $F/K$ , and put  $G = \text{Gal}(F/K)$ .

Suppose that  $-1$  is not a norm of from  $F$  to  $K$  of any unit but  $-1 \in N_G(\mathcal{O}_F^\times) \mathcal{O}_F^{\times 2}$ . First,  $N_G(\mathcal{O}_F^\times) \subset \{1, \pm \varepsilon\}$  modulo squares. The equations  $-1 = z^2$  and  $-1 = \varepsilon z^2$  have no solutions with  $z \in \mathcal{O}_F^\times$  for sign reasons. Hence the only possible solution is that  $-1 = -\varepsilon z^2$ , and then, necessarily  $F = K(\sqrt{\varepsilon})$ .

Suppose now that  $\varepsilon$  is not a norm of any unit in  $F/K$ . As before, if we test the condition  $\varepsilon \in N_G(\mathcal{O}_F^\times) \mathcal{O}_F^{\times 2}$ , we see the equations  $\varepsilon = -z^2$ , and  $\varepsilon = -\varepsilon^a z^2$  have no solution for sign reasons. Suppose that  $\varepsilon = \varepsilon^a z^2$  for

some odd integer  $a$  with  $\varepsilon^a \in N_G(\mathcal{O}_F^\times)$ . As  $N_G(\varepsilon) = \varepsilon^2$ , it is easy to see this implies  $\varepsilon \in N_G(\mathcal{O}_F^\times)$ , which contradicts our assumption. Thus  $a$  is even, and we conclude that  $\varepsilon \in \mathcal{O}_F^{\times 2}$ , i.e.  $F = K(\sqrt{\varepsilon})$ .

Hence, in any quadratic subextension  $F/K$  in  $K_\theta/K$  such that  $F \neq K(\sqrt{\varepsilon})$ , one has that the map  $N_G : \mathcal{E}_F \rightarrow \frac{\mathcal{O}_K^\times}{\mathcal{O}_K^\times \cap \mathcal{O}_F^{\times 2}}$  is not onto, and the result follows as in the imaginary case.  $\square$

### 3. Applications

Throughout this section, we explore applications of the previous results, including:

- How  $\lambda$  and deficiency change as we move up the tower  $K_\theta/K$ ;
- That  $\text{Def}(G_\theta) = 0$  implies the same for open subgroups of  $G_\theta$  when  $\delta = 1$ ;
- The rapid growth of  $\lambda$  as we move up a  $p$ -adic analytic quotient tower of  $G_\theta$ . The Tame Fontaine–Mazur conjecture predicts that infinite  $p$ -adic analytic quotients of  $G_\theta$  do not exist; thus, proving  $\lambda$  cannot grow rapidly would lend support to the Fontaine–Mazur conjecture;
- Some results in the direction of better understanding the cohomological dimension of  $G_\theta$ ;
- A computable test for maximality of  $\text{Def}(G_\theta)$ .
- We recall a well-known example of a quadratic imaginary field  $K$  where, for  $p = 3$ , we have  $r(G_{K,\theta}) = d(G_{K,\theta}) = 3$  so  $\text{Def}(G_{K,\theta}) = 0$  and  $|G_{K,\theta}| = \infty$ . We give examples of degree 8 extensions  $M/K$  whose 3-Hilbert class field tower is the base change of  $G_{K,\theta}$ , thus providing examples of fields  $M$  where  $|G_{M,\theta}| = \infty$  and  $\text{Def}(G_{M,\theta}) = 0$ . The infinite towers over  $M$  have 7 independent Minkowski units.

#### 3.1. Conserving the deficiency along the tower

Let  $F$  be a number field in the tower  $K_\theta/K$  and recall that  $F_\theta = K_\theta$ . We denote by  $\lambda_{F_\theta/F}$  the asymptotic Minkowski rank in  $F_\theta/F$ .

**PROPOSITION 3.1.** — *One has  $\lambda_{F_\theta/F} \geq [F : K]\lambda_{K_\theta/K}$ .*

*Proof.* — Let  $L \supset F \supset K$  in  $K_\theta/K$  be a large enough number field so that  $\lambda_{L/K} = \lambda_{K_\theta/K}$  and  $\lambda_{L/F} = \lambda_{F_\theta/F}$ . Set  $G = \text{Gal}(L/K)$  and  $H = \text{Gal}(L/F)$ . Then  $\mathcal{E}_L = \mathbb{F}_p[G]^{\lambda_{K_\theta/K}} \oplus N$ , where  $N$  is  $G$ -torsion. The result follows by noting that  $\mathbb{F}_p[G] \simeq_H \mathbb{F}_p[H]^{[F:K]}$  (see Section 1.3.1).  $\square$



COROLLARY 3.2. — For every number field  $F$  in  $K_\emptyset/K$ , we have

$$\text{Def}(G_{F,\emptyset}) \leq d(\mathcal{O}_F^\times) - [F : K]\lambda_{K_\emptyset/K}.$$

*Proof.* — This follows immediately from Theorem 2.9 and Proposition 3.1.  $\square$

*Remark 3.3.* — When  $\delta = 0$  the above Corollary is a consequence of strictly group-theoretic considerations. Namely, from equations (5.2) and (5.4) of [15] one deduces that for an open subgroup  $H$  of a pro- $p$  group  $G$ , one has

$$\text{Def}(H) + 1 \leq (G : H)(\text{Def}(G) + 1).$$

### 3.2. When $\text{Def}(G_\emptyset) = 0$

COROLLARY 3.4. — Let  $K$  be a number field containing  $\zeta_p$ . Suppose that  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\emptyset}^\times)^p = (\mathcal{O}_K^\times)^p$ , and that  $\text{Def}(G_\emptyset) = 0$ . Then, for every finite extension  $F/K$  in  $K_\emptyset/K$ , one has  $\text{Def}(G_{F,\emptyset}) = 0$ .

*Proof.* — Applying Theorem 2.9, we see  $\lambda = d(\mathcal{O}_K^\times)$  and is maximal and hence constant in the tower  $K_\emptyset/K$ , relative to the base field  $K$ . By Proposition 3.1 we see

$$\lambda_{F_\emptyset/F} \geq [F : K]\lambda = [F : K]d(\mathcal{O}_K^\times) = d(\mathcal{O}_F^\times).$$

The result follows by Theorem 2.9.  $\square$

COROLLARY 3.5. — Let  $G$  be a pro-2 group such that:

- (i)  $\text{Def}(G) = 0$ ,
- (ii) there exists a normal open subgroup  $H$  of  $G$  such that  $r(H) \neq d(H)$ .

Then  $G$  cannot be realized as the 2-tower of an imaginary quadratic field  $K$  of discriminant  $\text{disc}_K \equiv 1 \pmod{4}$  nor  $\text{disc}_K \equiv 0 \pmod{8}$ .

*Proof.* — The discriminant hypotheses imply  $-1 \notin \mathcal{O}_{K_\emptyset}^{\times 2}$ . The result follows from Corollary 3.4.  $\square$

The condition that the number of Minkowski units is maximal is very strong<sup>(2)</sup>:

PROPOSITION 3.6. — The finite  $p$ -groups  $G$  that have the property  $\text{Def}(H) = 0$  for every subgroup  $H$  of  $G$  are exactly the cyclic groups and generalized quaternion group  $Q_{2^n} = \langle x, y \mid x^{2^{n-1}} = 1, x^{2^{n-2}} = y^2, yxy^{-1} = x^{-1} \rangle$ ,  $n \geq 3$ .

<sup>(2)</sup> We thank Ozaki for bringing this result to our attention.

*Proof.* — For  $G$  with this property, every abelian subgroup  $H$  of  $G$  is of deficiency 0, forcing  $H$  to be cyclic. Then,  $G$  is cyclic or the generalized quaternion group  $Q_{2^n}$  of order  $2^n$  (see for example [34, Theorem 9.7.3]). For the converse, obviously cyclic groups  $G$  satisfy  $\text{Def}(H) = 0$  for every subgroup  $H$  of  $G$ . Concerning  $Q_{2^n}$ , recall that its subgroups are cyclic or isomorphic to  $Q_{2^{n-1}}$ , and that the Schur multipliers of the generalized quaternion groups  $Q_{2^k}$  are trivial (or in other words that  $\text{Def}(Q_{2^k}) = 0$ ).  $\square$

*Remark 3.7.* — Take  $p = 2$ , and let  $K$  be an imaginary quadratic field. Recall that  $\text{Def}(G_\emptyset) \in \{0, 1\}$ . We suspect that when  $G_\emptyset$  is infinite then  $\text{Def}(G_\emptyset)$  is maximal. If this is not the case and the hypothesis of Corollary 3.4 holds ( $\text{disc}_K \equiv 1 \pmod{4}$  or  $\text{disc}_K \equiv 0 \pmod{8}$ ), then  $r(H) = d(H)$  for every open normal subgroup  $H$  of  $G_\emptyset$ .

*Remark 3.8.* — Observe that Poincaré pro- $p$  groups of dimension 3 are examples of infinite groups satisfying the condition of Corollary 3.4, see for example [24, Chapter III, Section 7].

We close this subsection with an explicit, albeit contrived, example with  $p = 2$ .

*Example 3.9.* — Let  $K = \mathbb{Q}(\sqrt{-3 \cdot 5 \cdot 53})$ . An easy MAGMA computation gives that the class group of  $K$  is  $(2, 2)$  and its 2-Hilbert class field tower has degree 8 over  $K$ . Straightforward computations show this group has at least three cyclic subgroups of order 4, hence it is the quaternion group of order 8. Here  $\mathcal{O}_K^\times = \{1, -1\}$ , and as the discriminant of  $K$  is prime to 4,  $i^2 = -1 \notin \mathcal{O}_{K_0}^{\times 2}$  so  $\mathcal{O}_K^\times \cap \mathcal{O}_{K_0}^{\times 2} = \{1\}$  and  $\beta = 0$ . Then Theorem 2.9 gives  $\text{Def}(G_\emptyset) = 1 - \lambda$ . But it is well-known the quaternion group has deficiency 0 so  $\lambda = 1$ . There is a Minkowski unit in this (short) tower. Indeed, if one computed a basis of the units of the degree 16 field that is the top of the tower and computed norms to  $K$ , the elements with norm  $-1$  (which exist!) are Minkowski units. We have not explored computing the Minkowski unit explicitly in this case, but knowledge of the unit group of the genus field should make such an effective calculation possible.

### 3.3. In the context of the Fontaine–Mazur conjecture

The conjecture of Fontaine–Mazur [7, Conjecture 5a] asserts that every analytic quotient of  $G_\emptyset$  must be finite. By class field theory, one knows that every infinite analytic quotient of  $G_\emptyset$  must be of analytic dimension at least 3 (see [20, Proposition 2.12]).

One knows that  $G_\emptyset$  is not  $p$ -analytic when the  $p$ -rank  $d(\text{Cl}_K)$  of the class group  $\text{Cl}_K$  of  $K$  is large compared to  $[K : \mathbb{Q}]$ . See A.3.11 of [18]. Alternatively, this is (literally!) an exercise on page 78 of [35].

Suppose  $G := G_\emptyset$  is infinite and analytic. One knows that every infinite analytic pro- $p$  group contains an open *uniform* subgroup. To simplify, assume  $G$  is uniform. Denote by  $(G_n)$  the  $p$ -central descending series of  $G$  (it is also the Frattini series), and let  $K_n = K_\emptyset^{G_n}$ . Put  $H_n = \text{Gal}(K_n/K)$ ; recall that  $\#H_n = p^{dn}$ , where  $d = d(G_\emptyset)$  is also the dimension of  $G_\emptyset$  as analytic group. For  $n \geq 1$ , denote by  $\lambda_n$  the Minkowski-rank of the units along  $K_\emptyset/K_n$ .

The hypothesis of Corollary 3.10 below is, assuming the Fontaine–Mazur Conjecture, never satisfied. We include the Corollary to indicate a possible strategy to prove  $G_\emptyset$  is not analytic, namely show the number of Minkowski units does not grow so rapidly in the tower.

**COROLLARY 3.10.** — *Let  $G_\emptyset$  be pro- $p$  analytic of dimension  $d$ . Then for  $m$  large,*

$$\begin{aligned} (r_1 + r_2)[K_m : K] - 1 + \delta - \frac{d(d-1)}{2} &\leq \lambda_m \\ &\leq (r_1 + r_2)[K_m : K] - 1 + \delta - \frac{d(d-3)}{2}. \end{aligned}$$

*Proof.* — Theorem 2.9 here, Theorem 4.35 of [6], and the assumption that  $G_\emptyset$  is uniform imply  $\text{Def}(G_m)$  is constant and equal to  $\text{Def}(G_\emptyset) = \frac{d(d-3)}{2}$ . As remarked in the introduction,  $\beta_m \leq d(G_m) = d(G) = d$ . We immediately see  $\lambda_m \sim (r_1 + r_2)[K_m : K]$ , proving the main terms in the estimates.

We now prove the more refined estimates. Let us choose  $n \gg m \gg 0$  such that:

$$\mathcal{E}_{K_n} = \mathbb{F}_p[H_{n,m}]^{\lambda_m} \oplus N_{n,m} = \mathbb{F}_p[H_n]^\lambda \oplus N_n$$

where  $\lambda = \lambda_{K_\emptyset/K}$ ,  $\lambda_m = \lambda_{K_\emptyset/K_m}$ ,  $H_{n,m} = \text{Gal}(K_n/K_m)$ , and  $N_{n,m}$  and  $N_n$  are torsion modules over  $\mathbb{F}_p[H_{n,m}]$  and  $\mathbb{F}_p[H_n]$  respectively.

Then by Proposition 3.1, we see  $\lambda_m = [K_m : K]\lambda + \lambda_{m,n}^{\text{new}}$ ; the quantity  $\lambda_{m,n}^{\text{new}}$  corresponds to the  $\mathbb{F}_p[H_{n,m}]$ -free part in  $N_n$ . Hence, by Theorem 2.9, one has

$$\text{Def}(G_m) \geq d(\mathcal{O}_{K_m}^\times) - \lambda_m - \beta_m \geq d(\mathcal{O}_{K_m}^\times) - [K_m : K]\lambda - \lambda_{m,n}^{\text{new}} - d(G_m).$$

After noting that  $d(\mathcal{O}_{K_m}^\times) = (r_1 + r_2)[K_m : K] - 1 + \delta$ , we get

$$(3.1) \quad \text{Def}(G_m) \geq (r_1 + r_2 - \lambda)[K_m : K] - 1 + \delta - d - \lambda_{m,n}^{\text{new}}$$

But  $\text{Def}(G_m) = \text{Def}(G_\emptyset) = \frac{d(d-3)}{2}$ . Hence (3.1) becomes

$$(3.2) \quad \lambda_{m,n}^{\text{new}} \geq (r_1 + r_2 - \lambda)[K_m : K] - 1 + \delta - \frac{d(d-1)}{2}.$$

and

$$(3.3) \quad \lambda_m \geq (r_1 + r_2)[K_m : K] - 1 + \delta - \frac{d(d-1)}{2},$$

proving the first inequality. The upper bound follows as  $\text{Def}(G_m) \leq d(\mathcal{O}_{K_m}^\times) - \lambda_m$  so

$$(3.4) \quad \lambda_m \leq (r_1 + r_2)[K_m : K] - 1 + \delta - \frac{d(d-3)}{2}. \quad \square$$

### 3.4. On the cohomological dimension of $G_\emptyset$

Since the works of Labute [16], Labute–Mináč [17] and Schmidt [31], etc. one knows that in certain cases the groups  $G_S$ , for  $S$  tame, are of cohomological dimension 2. In all the examples of these papers  $S \neq \emptyset$ . The question of the computation of cohomological dimension of  $G_\emptyset$  is still an open problem (one can find partial negative answers in [21]). To prove Theorem 3.12, we need the following lemma due to Schmidt [29, Proposition 1]. We first need to define the level 3 partial Euler characteristic of a pro- $p$  group  $G$  as

$$\chi_3(G) := 1 - d(G) + r(G) - \dim H^3(G) = 1 + \text{Def}(G) - \dim H^3(G).$$

LEMMA 3.11 (Schmidt). — *Let  $G$  be an infinite pro- $p$  group such that for a fixed constant  $n \geq 0$  and every open subgroup  $H$  of  $G$ , one has*

$$-\chi_3(H) + n \geq -[G : H]\chi_3(G).$$

Then  $\text{cd}(G) \leq 3$ .

THEOREM 3.12. — *Let  $K$  be a number field such that*

- (i)  $K$  contains a primitive  $p$ th root of unity;
- (ii)  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\emptyset}^\times)^p = (\mathcal{O}_K^\times)^p$ .

Then  $\dim H^3(G_\emptyset) > 0$ . Moreover:

- If  $\dim H^3(G_\emptyset) = 1$ , then  $G_\emptyset$  is finite or of cohomological dimension 3;
- If  $\text{Def}(G_\emptyset) = 0$ , and if  $G_\emptyset$  is of cohomological dimension 3, then  $G_\emptyset$  is a Poincaré duality group.

*Proof.* — As  $\mathcal{O}_K^\times \cap (\mathcal{O}_{K_\theta}^\times)^p = (\mathcal{O}_K^\times)^p$  and  $\delta = 1$ , one has, by Theorem 2.9,

$$\text{Def}(G_\theta) = d(\mathcal{O}_K^\times) - \lambda_{K_\theta/K} = r_1 + r_2 - \lambda_{K_\theta/K}.$$

Let  $H$  be an open normal subgroup of  $G_\theta$  and set  $F = K_\theta^H$ . Proposition 3.1 implies  $\lambda_{F_\theta/F} \geq \lambda_{K_\theta/K}[G_\theta : H]$ , so Theorem 2.9 implies

$$\text{Def}(H) \leq [G_\theta : H](r_1 + r_2 - \lambda_{K_\theta/K}).$$

Recalling that  $\chi_2$  is the Euler characteristic truncated at second cohomology,

$$\chi_2(H) = 1 + \text{Def}(H) \leq 1 + [G_\theta : H](r_1 + r_2 - \lambda_{K_\theta/K}),$$

so  $\chi_2(H)$  cannot be equal to  $[G_\theta : H]\chi_2(G_\theta) = [G_\theta : H](1 + r_1 + r_2 - \lambda_{K_\theta/K})$ , a necessary condition, by Theorem 5.4 of [15], for  $G_\theta$  to be of cohomological dimension 2. Hence  $G_\theta$  is not of cohomological dimension 2 so  $\dim H^3(G_\theta) > 0$ .

Now suppose  $G_\theta$  is infinite and  $\dim H^3(G_\theta) = 1$ . By Theorem 2.9 and Proposition 3.1, one has

$$\begin{aligned} [-1 - \text{Def}(H) + \dim H^3(H)] + 1 &= \lambda_{F_\theta/F} - d(\mathcal{O}_F^\times) + \dim H^3(H) \\ &\geq [G_\theta : H](\lambda_{K_\theta/K} - (r_1 + r_2)) \\ &= [G_\theta : H](-1 - \text{Def}(G_\theta) + \dim H^3(G_\theta)) \end{aligned}$$

where the last equality follows from Theorem 2.9 using that  $\beta = 0$  and  $\dim H^3(G_\theta) = 1$ . Now take  $n = 1$  in Lemma 3.11 to conclude  $\text{cd}(G_\theta) = 3$ .

Finally, to check that our group is a Poincaré group, following [24, Chapter III, Section 7], we need only verify that  $D_i(\mathbb{Z}/p) := \lim_{\substack{\rightarrow \\ U}} H^i(U)^\wedge = 0$  for  $i = 0, 1, 2$ , where the limit is taken over open subgroups  $U$  of  $G_\theta$  and the transition maps are dual to the corestriction. Recall that cohomological dimension is nonincreasing when one restricts to a closed subgroup and that cyclic groups have infinite cohomological dimension, so as  $G_\theta$  is assumed to be of finite cohomological dimension, it is infinite and thus  $D_0(\mathbb{Z}/p) = 0$ . Moreover that

$$D_1(\mathbb{Z}/p) = \lim_{\substack{\rightarrow \\ U}} U^{ab}/p = 0$$

follows from the proof of the Principal Ideal Theorem: Namely, for a group  $G$  let  $G'$  be its (closed) commutator subgroup and let  $G''$  be the (closed) commutator subgroup of  $G'$ . The key part of the proof of the Principal Ideal Theorem is that the transfer map

$$\text{Ver} : G/G' \longrightarrow G'/G''$$

is the zero map. As the transfer map is the dual of the corestriction map,  $D_1(\mathbb{Z}/p) = 0$ .

We now show  $D_2(\mathbb{Z}/p) = 0$ . For  $U \subset G_\emptyset$  open, taking the  $U$ -cohomology of the short exact sequence of trivial  $U$ -modules

$$0 \longrightarrow \mathbb{Z}/p \longrightarrow \mathbb{Q}/\mathbb{Z} \xrightarrow{p} \mathbb{Q}/\mathbb{Z} \longrightarrow 0$$

gives

$$H^1(U, \mathbb{Q}/\mathbb{Z}) \xrightarrow{p} H^1(U, \mathbb{Q}/\mathbb{Z}) \longrightarrow H^2(U, \mathbb{Z}/p)$$

which yields  $(U^{ab})^\vee/p \hookrightarrow H^2(U, \mathbb{Z}/p)$ . As  $\dim(U^{ab})^\vee/p = \dim H^1(U)$  and  $\text{Def}(U) = 0$  by Corollary 3.4, this injection is an isomorphism and  $H^2(U, \mathbb{Z}/p)^\wedge \simeq U^{ab}/p \simeq H^1(U, \mathbb{Z}/p)^\wedge$ . Since the duals of the two corestriction maps are induced by the transfer,  $D_1(\mathbb{Z}/p) = 0 \implies D_2(\mathbb{Z}/p) = 0$ .  $\square$

*Remark 3.13.* — The first part of Theorem 3.12 extends the following observation that can be deduced from the relationship between Galois cohomology and étale cohomology.

We use the formalism of étale cohomology as in [23]. Suppose  $\text{Def}(G_\emptyset)$  is maximal. Then the étale version of the Hochschild–Serre spectral sequence with [30, Theorem 3.4] shows that  $H^i(G_\emptyset) \simeq H_{\text{ét}}^i(\text{Spec } \mathcal{O}_K)$  for  $i = 1, 2$ . Moreover, if  $G_\emptyset$  has cohomological dimension 2, then  $G_\emptyset$  is infinite: by [30, Lemma 3.7]) and from the Hochschild–Serre spectral sequence we also get  $\{0\} = H^3(G_\emptyset) \simeq H_{\text{ét}}^3(\text{Spec } \mathcal{O}_K) \simeq \mu_{K,p}$ , where here  $\mu_{K,p} = \langle \zeta_p \rangle \cap K$  (by [30, Theorem 3.4]). Hence  $\delta$  must be zero.

### 3.5. Detecting maximality

The strategy of the Hochschild–Serre spectral sequence allows us to prove Theorem 3.14 below, a computationally feasible method of showing  $\text{III}_\emptyset^2$  is maximal.

**THEOREM 3.14.** — *Suppose there exist two linearly disjoint unramified  $\mathbb{Z}/p$ -extensions  $F_1/K$  and  $F_2/K$  such that  $t_{G_i}(\mathcal{E}_{F_i}) = 0$ ,  $i = 1, 2$ , where  $G_i = \text{Gal}(F_i/K)$ . Then  $\text{Def}(G_\emptyset) = d(\mathcal{O}_K^\times)$  is maximal. Only one such extension  $F_i/K$  is sufficient if  $F_i \not\subset K'(\sqrt[p]{\mathcal{O}_K^\times})$ , which is the case when  $\delta = 0$ .*

*Proof.* — We use the notations of Section 2.2. First note that Lemma 2.6 and the fact that  $t_{G_i}(\mathcal{E}_{F_i}) = 0$  implies  $\lambda = 0$ .

If  $\delta = 0$ , Proposition 2.8 implies  $\beta = 0$  so  $\text{Def}(G_\emptyset)$  is maximal by Theorem 2.9.

We now address the  $\delta = 1$  case. First suppose  $F_1 \not\subset K(\sqrt[p]{\mathcal{O}_K^\times})$ . Then one can choose a set  $S_2$  consisting of  $d(\mathcal{O}_K^\times)$  primes  $\mathfrak{p}$  of  $K$  that split completely in  $F_1$  and whose Frobenius automorphisms form a basis of

$\text{Gal}(K(\sqrt[p]{\mathcal{O}_K^\times})/K)$ . Since  $t_{G_1}(\mathcal{E}_{F_1}) = 0$ , by Theorem 1.1 we see that for each  $\mathfrak{p} \in S_2$  there is a  $\mathbb{Z}/p$ -extension of  $F_1$ , and hence of  $K_\theta$ , ramified only at (the primes above)  $\mathfrak{p}$ . Each of these elements gives rise to a relation of  $G_\theta$ . As usual, one gets the rest of the relations “for free” by choosing primes that split completely in  $K(\sqrt[p]{\mathcal{O}_K^\times})/K$  but form a basis of  $\text{Gal}(K(\sqrt[p]{\mathcal{O}_K^\times})/K)$ . For such primes  $\mathfrak{p}$  there is always an abelian extension of  $K$  ramified only at  $\mathfrak{p}$ , also giving rise to a relation of  $G_\theta$ .

We study the remaining case, namely when  $F_1, F_2 \subset K(\sqrt[p]{\mathcal{O}_K^\times})$ . Choose a prime  $\mathfrak{q}_1$  of  $K$  such that its Frobenius generates  $\text{Gal}(F_1/K)$  and  $\mathfrak{q}_1$  splits in  $F_2$ . Choose  $\mathfrak{q}_2$  similarly. Then, as before, when we allow ramification at  $\mathfrak{q}_1$  we obtain a ramified extension over  $F_{2,\theta}$  and when we allow ramification at  $\mathfrak{q}_2$  we obtain a ramified extension over  $F_{1,\theta}$ . In this case we build  $S_2$  by starting with  $S_2 = \{\mathfrak{q}_1, \mathfrak{q}_2\}$  and augmenting it to include primes that split completely in  $F_1F_2$  and whose Frobenius automorphisms, along with those of  $\mathfrak{q}_1$  and  $\mathfrak{q}_2$ , form a basis of  $\text{Gal}(K(\sqrt[p]{\mathcal{O}_K^\times})/K)$ . For each of these primes when we allow ramification at  $\mathfrak{p}$  we obtain a ramified extension over  $F_{i,\theta}$  for  $i = 1, 2$ . Each of these primes gives rise to a relation of  $G_\theta$  and along with the “free relations” we get  $\text{Def}(G_\theta) = d(\mathcal{O}_K^\times)$ .  $\square$

### 3.6. Infinite towers having Minkowski units

In this section, as we compare towers of different fields, we include the field in the subscript of  $G$ . We produce examples of number fields  $K$  for which  $G_{K,\theta}$  is infinite,  $\beta = 0$ , and  $\text{Def}(G_{K,\theta})$  is minimal. By Theorem 2.9  $\lambda > 0$  in these cases. We first need some set up.

#### 3.6.1. Stability of towers

Let us start with the following context. Let  $K/k$  be a Galois extension of order coprime to  $p$ . Observe that  $K_\theta/k$  is Galois. Put  $\Delta := \text{Gal}(K/k)$  and  $\Gamma = \text{Gal}(K_\theta/k)$ . By the Schur–Zassenhaus Theorem one has  $\Gamma \simeq G_{K,\theta} \rtimes \Delta$ .

Obviously  $\text{Gal}(Kk_\theta/K) \simeq G_{k,\theta}$  and there is a natural surjective map  $G_{K,\theta} \twoheadrightarrow G_{k,\theta}$ . In [11] Gras shows:

**THEOREM 3.15.** — *Suppose  $K/k$  is Galois and of degree prime to  $p$ . Then  $G_{K,\theta} \simeq G_{k,\theta}$  if and only if  $d(G_K) = d(G_k)$ .*

We give an alternative proof of this result inspired by Wingberg [39, 41].

First, let  $N$  be the closed normal subgroup of  $G_{K,\theta}$  generated by the  $g^s g^{-1}, g \in G_{K,\theta}, s \in \Delta$ ; put  $G_{K,\theta}^+ := G_{K,\theta}/N$ . In other words,  $G_{K,\theta}^+$  is the

largest quotient of  $G_{K,\emptyset}$  on which  $\Delta$  acts trivially. Since  $\Delta$  acts trivially on  $G_{k,\emptyset}$ , the morphism  $\varphi : G_{K,\emptyset} \rightarrow G_{k,\emptyset}$  factors through  $G_{K,\emptyset}^+$ .

LEMMA 3.16. — *One has  $H^1(G_{k,\emptyset}) \simeq H^1(G_{K,\emptyset}^+)$ .*

*Proof.* — By base change from  $k$  to  $K$ , using that  $(\#\Delta, p) = 1$ , we have an injection  $H^1(G_{k,\emptyset}) \hookrightarrow H^1(G_{K,\emptyset}^+)$ . As elements of the  $H^1(G_{K,\emptyset}^+)$  correspond to  $\mathbb{Z}/p$ -extensions of  $K$  on whose Galois group  $\Delta$  acts trivially, these descend to  $k$ , so the injection is a surjection.  $\square$

Consider now  $\Gamma(p)$  the maximal pro- $p$  quotient of  $\Gamma$ .

LEMMA 3.17. — *One has  $\Gamma(p) \simeq G_{k,\emptyset}$ .*

*Proof.* — Obviously,  $\Gamma(p) \twoheadrightarrow G_{k,\emptyset}$ . But it is easy to see that  $\Gamma(p)$  corresponds to a pro- $p$  extension of  $k$  unramified everywhere. By maximality, we deduce the result.  $\square$

We recall now the result of Wingberg (given in [39] when  $\Delta \simeq \mathbb{Z}/2$ , and in general in an unpublished work [41]).

PROPOSITION 3.18 (Wingberg). — *If  $d(G_{K,\emptyset}) = d(G_{k,\emptyset})$  then the natural map  $G_{K,\emptyset}^+ \twoheadrightarrow G_{k,\emptyset}$  is an isomorphism and induces the injection  $H^2(G_{K,\emptyset}^+) \hookrightarrow H^2(G_{K,\emptyset})^\Delta$ .*

*Proof.* — The Hochschild–Serre spectral sequence for

$$1 \longrightarrow M \longrightarrow G_{K,\emptyset}^+ \longrightarrow G_{k,\emptyset} \longrightarrow 1$$

gives the commutative diagram (by Lemmas 3.16 and 3.17):

$$\begin{array}{ccccc}
 0 & \longrightarrow & H^1(M)^{G_{k,\emptyset}} & \longrightarrow & H^2(G_{k,\emptyset}) & \longrightarrow & H^2(G_{K,\emptyset}^+) \\
 & & & & \downarrow \simeq & & \downarrow \text{Inf} \\
 & & & & H^2(\Gamma(p)) & & \\
 & & & & \downarrow \text{Inf} & & \\
 & & & & H^2(\Gamma) & \hookrightarrow & H^2(G_{K,\emptyset})^\Delta
 \end{array}$$

We then deduce that  $H^1(M)^{G_{k,\emptyset}} = 0$ , or equivalently that  $M = \{e\}$  so  $G_{K,\emptyset}^+ \twoheadrightarrow G_{k,\emptyset}$  is an isomorphism and  $H^2(G_{k,\emptyset}) \simeq H^2(G_{K,\emptyset}^+) \hookrightarrow H^2(G_{K,\emptyset})^\Delta$ .  $\square$

We can now give an alternative proof of Theorem 3.15.

*Proof.* — Take the Hochschild–Serre spectral sequence of

$$1 \longrightarrow N \longrightarrow G_{K,\emptyset} \longrightarrow G_{K,\emptyset}^+ \longrightarrow 1$$



to obtain

$$0 \longrightarrow H^1(G_{K,\emptyset}^+) \longrightarrow H^1(G_{K,\emptyset}) \longrightarrow H^1(N)^{G_{K,\emptyset}^+} \longrightarrow H^2(G_{K,\emptyset}^+) \longrightarrow H^2(G_{K,\emptyset}).$$

By Proposition 3.18, one gets:

$$0 \longrightarrow H^1(G_{K,\emptyset}^+) \longrightarrow H^1(G_{K,\emptyset}) \longrightarrow H^1(N)^{G_{K,\emptyset}^+} \longrightarrow 0.$$

Recall that  $N = \{e\}$  if and only if  $H^1(N)^{G_{K,\emptyset}^+} = 0$ . Hence  $G_{K,\emptyset} \simeq G_{K,\emptyset}^+ \simeq G_{k,\emptyset}$  if and only if  $d(G_{K,\emptyset}^+) = d(G_{K,\emptyset})$ , if and only if  $d(G_{k,\emptyset}) = d(G_{K,\emptyset})$ .  $\square$

### 3.6.2. Example

Let  $k = \mathbb{Q}(\sqrt{-3321607})$  and  $p = 3$ . It is easy to compute that  $d = r = 3$ . Furthermore, the  $\text{Gal}(k/\mathbb{Q})$  action on  $G_{k,\emptyset}$  gives that the depth of each relation is at least three so we may take  $1 - 3t + 3t^3$  as a Golod–Shafarevich polynomial for  $G_k$ . As this polynomial has a root in  $]0, 1[$  we see  $|G_{k,\emptyset}| = \infty$ . This example is well-known (see [33]).

We will find explicit multi-quadratic fields  $M/\mathbb{Q}$  such that  $d(G_{Mk,\emptyset}) = d(G_{k,\emptyset}) = 3$ . Theorem 3.15 then gives  $(Mk)_\emptyset = M(k_\emptyset)$ . The largest  $M$  we give is degree 8 so  $Mk$  is a totally complex field of degree 16 with  $r = d = 3$  and  $\text{Def}(G_{Mk,\emptyset}) = 0$ . In this case  $\lambda = 7$ , that is there are 7 independent Minkowski units all the way up the tower  $(Mk)_\emptyset/Mk$ .

**LEMMA 3.19.** — *For  $M/\mathbb{Q}$  multi-quadratic, to check that  $d(G_{Mk,\emptyset}) = d(G_{k,\emptyset}) = 3$ , it is equivalent to check that  $d(G_{Nk,\emptyset}) = d(G_{k,\emptyset}) = 3$  for every quadratic extension  $N/k$  contained in  $Mk$ .*

*Proof.* — Suppose  $d(G_{Mk,\emptyset}) > 3$ . The elementary 2-abelian group  $\text{Gal}(Mk/k)$  acts on the  $\mathbb{F}_3$ -vector space  $G_{Mk,\emptyset}^{3-el,ab}$ . Since 2 and 3 are relatively prime and both square roots of unity lie in  $\mathbb{F}_3$ ,  $G_{Mk,\emptyset}^{3-el,ab}$  decomposes as a direct sum of one-dimensional  $\mathbb{F}_3[\text{Gal}(Mk/k)]$  subspaces. The trivial summands descend to  $k$ . Since we are assuming  $d(G_{Mk,\emptyset}) > 3$ , there is a non-trivial summand. The kernel of the action on this summand is  $\text{Gal}(Mk/N)$  where  $[N : k] = 2$ . Thus any extra generators of  $G_{Mk,\emptyset}$  are realized over some  $N$ .  $\square$

**PROPOSITION 3.20.** — *For  $M = \mathbb{Q}(\sqrt{3}, \sqrt{7}, \sqrt{337})$  we have  $|G_{Mk,\emptyset}| = \infty$ ,  $d(G_{Mk,\emptyset}) = d(G_{k,\emptyset}) = 3$  and  $\text{Def}(G_{Mk,\emptyset}) = 0$  is minimal. Assuming the GRH, for  $M = \mathbb{Q}(\sqrt{3}, \sqrt{7}, \sqrt{r})$  with  $r = 383$  or  $r = 461$  we have  $|G_{Mk,\emptyset}| = \infty$ ,  $d(G_{Mk,\emptyset}) = d(G_{k,\emptyset}) = 3$  and  $\text{Def}(G_{Mk,\emptyset}) = 0$  is minimal.*

*Proof.* — As  $|G_{k,\emptyset}| = \infty$ , we immediately have  $|G_{Mk,\emptyset}| = \infty$ . We used GP-PARI and MAGMA to check  $d(G_{N,\emptyset}) = 3$  for the seven quadratic extensions  $N/k$  inside  $Mk/k$ . For the computations with  $r = 337$ , we did not assume the GRH and these took several days. For  $r = 383$  and  $461$  we assumed the GRH and the computations took several minutes. Theorem 3.15 implies  $(Mk)_\emptyset = M(k_\emptyset)$  so  $r(\text{Gal}((Mk)_\emptyset/Mk)) = r(\text{Gal}(k_\emptyset/k)) = 3$  which in turn gives  $\text{Def}(G_{Mk,\emptyset}) = 7$ .  $\square$

*Remark 3.21.* — The fields  $M$  above were found by checking all primes  $q$  less than 500 such that  $d(G_{k(\sqrt{q}),\emptyset}) = d(G_{k,\emptyset}) = 3$ . The first such prime was 3. We then searched this list for a second prime  $q_2$  such that for every quadratic extension  $N/k$  inside  $k(\sqrt{3}, \sqrt{q_2})$  we had  $d(G_{N,\emptyset}) = 3$ . We found the first  $q_2 = 7$ . Finally, we searched for  $q_3$  such that for every quadratic extension  $N/k$  inside  $k(\sqrt{3}, \sqrt{7}, \sqrt{q_3})$  we had  $d(G_{N,\emptyset}) = 3$ .

The natural question arises as to whether there are infinitely many fields  $M$  with  $M/k$  Galois of degree prime to 3 such that  $d(G_{Mk,\emptyset}) = 3$  so  $G_{Mk,\emptyset} = G_{k,\emptyset}$ . One may also ask: Does there exist a number field  $K$  such that  $|G_{K,\emptyset}| = \infty$  and  $\text{Def}(G_{K,\emptyset})$  is not maximal, but for every  $F \subset K$  we have either  $|G_{F,\emptyset}| < \infty$  or  $\text{Def}(G_{F,\emptyset})$  is maximal?

*Remark 3.22.* — Take  $p = 3$ . The previous approach does not allow us to produce situations with Minkowski units and such that  $\delta = 1$ . Indeed, let  $k = \mathbb{Q}(\sqrt{n})$  be a quadratic extension,  $n \in \mathbb{Z}$ ,  $n \notin \mathbb{Z}^2 \cup -3\mathbb{Z}^2$ , such that  $d = 3$ . By the “Spiegelungssatz” phenomenon of Scholz [32], the 3-rank of the class group of  $\mathbb{Q}(\sqrt{-3n})$  is at least 2, showing that  $k_\emptyset(\zeta_3) \neq (k(\zeta_3))_\emptyset$ .

#### 4. On the depth of the relations

In this section we show the existence of Minkowski units deep in the Frattini tower imply that some of the relations of  $G_\emptyset$  are very deep. This makes it “more likely” that one can prove  $G_\emptyset$  is infinite using the Golod–Shafarevich series. We also prove a converse, namely the existence of very deep relations implies the existence of Minkowski units along the Frattini tower. One reason we study the Frattini tower as opposed to the Zassenhaus tower is that it is easier to use software for computations along the Frattini tower of  $K_\emptyset/K$ .

## 4.1. On the Zassenhaus filtration

### 4.1.1. Basic properties

We refer to Lazard [18, Appendice A3]. Given a finitely presented pro- $p$  group  $G$ , let us take a minimal presentation of  $G$

$$1 \longrightarrow R \longrightarrow F \xrightarrow{\varphi} G \longrightarrow 1,$$

where  $F$  is a free pro- $p$  group on  $d$  generators; here  $d = d(G)$ . Let  $I = \ker(\mathbb{F}_p[[F]] \rightarrow \mathbb{F}_p)$  be the augmentation ideal of  $\mathbb{F}_p[[F]]$ , and for  $n \geq 1$  consider  $F^n = \{x \in F, x - 1 \in I^n\}$ . The sequence  $(F^n)$  of open subgroups of  $F$  is the Zassenhaus filtration of  $F$ .

The depth  $\omega$  of  $x \in F$  is defined as being  $\omega(x) = \max\{n, x - 1 \in I^n\}$ , with the convention that  $\omega(1) = \infty$ ; the function  $\omega$  is a valuation following terminology of Lazard. Hence  $F^n = \{g \in F, \omega(g) \geq n\}$ . This allows us to define a depth  $\omega_G$  on  $G$  as follows:  $\omega_G(x) = \max\{\omega(g), g \in F, \varphi(g) = x\}$ . Put  $G^n = \{x \in G, \omega_G(x) \geq n\}$ . Observe that  $G^n = F^n R / R \simeq F^n / (F^n \cap R)$ ; the sequence  $(G^n)$  is the Zassenhaus series of  $G$ , it corresponds to the filtration arising from the augmentation ideal  $I_G$  of  $\mathbb{F}[[G]]$ , see [18, Appendice A3, Theorem 3.5]. One has the following property. If  $\pi : G' \twoheadrightarrow G$  is surjective, then  $\omega_G$  is the restriction of  $\omega_{G'}$ ; in other words,

$$\omega_G(x) = \max\{\omega_{G'}(y), y \in G', \pi(y) = x\}.$$

Denote by  $(G_n)$  the Frattini filtration of  $G$ . Recall the well-known relationship between these two filtrations of  $G$ :

LEMMA 4.1. — *One has  $G_n \subset G^{2^{n-1}}$ .*

We say a few words about the reverse inclusions. Let  $H$  be an open normal subgroup of  $G$ . Since the groups  $(G^n)$  form a basis of neighborhoods of 1, let  $a(H)$  be the smallest integer such that  $G^{a(H)} \subset H$ . We want to give some estimates on  $a(H)$  in some special cases.

DEFINITION 4.2. — *For a pro- $p$  group  $\Gamma$ , denote by*

$$I_\Gamma := \ker(\mathbb{F}_p[[\Gamma]] \longrightarrow \mathbb{F}_p),$$

*the augmentation ideal of  $\mathbb{F}_p[[\Gamma]]$ ; and denote by  $k(\Gamma)$  the smallest integer such that  $I_\Gamma^{k(\Gamma)} = \{0\}$ , where we allow  $k(\Gamma) = \infty$ .*

PROPOSITION 4.3 ([15, Chapter 7, Section 7.6, Theorem 7.6]). — *Let  $1 \rightarrow H \rightarrow G \rightarrow G/H \rightarrow 1$  be an exact sequence of pro- $p$  groups. Then  $I_H = \ker(\mathbb{F}_p[[G]] \rightarrow \mathbb{F}_p[[G/H]])$ .*

Following Koch's book [15, Chapter 7, Section 7.4], we give some estimates for  $a(\mathbb{H})$ .

PROPOSITION 4.4. — *One has:*

- (i)  $a(\mathbb{H}) \leq k(\mathbb{G}/\mathbb{H}) \leq |\mathbb{G}/\mathbb{H}|$ .
- (ii) *If  $\Gamma' \triangleleft \Gamma$  are two finite  $p$ -groups, then  $k(\Gamma) \leq k(\Gamma/\Gamma')k(\Gamma')$ .*
- (iii)  $k(\mathbb{G}/\mathbb{G}_2) = p$ .

*Proof.*

(i). — Take  $k$  such that  $I_{\mathbb{G}/\mathbb{H}}^k = \{0\}$ . Then by Proposition 4.3 one has  $I_{\mathbb{G}}^k \subset I_{\mathbb{H}}$ , which implies  $\mathbb{G}^k \subset \mathbb{H}$ , and then  $a(\mathbb{H}) \leq k$ . In particular,  $a(\mathbb{H}) \leq k(\mathbb{G}/\mathbb{H})$ . For the second part of the inequality, observe that: for every finite  $p$ -group  $\Gamma$ , one has  $I_{\Gamma}^{|\Gamma|} = \{0\}$  (see the proof of Lemma 7.4 of [15, Chapter 7, Section 7.4]), showing that  $k(\Gamma) \leq |\Gamma|$ .

(ii). — By Proposition 4.3, one has  $I_{\Gamma}^{k(\Gamma/\Gamma')} \subset I_{\Gamma'}$ , and then  $I_{\Gamma}^{k(\Gamma/\Gamma')k(\Gamma')} \subset I_{\Gamma'}^{k(\Gamma')} = \{0\}$ .

(iii). — This follows as  $\mathbb{G}/\mathbb{G}_2$  is  $p$ -elementary abelian. □

For every integer  $n \geq 1$ , put  $a_n := a(\mathbb{G}_{n+1})$ . Observe that  $a_1 = 1$ .

PROPOSITION 4.5. — *For  $n \geq 2$ , one has  $a_n \leq p^n$ . Therefore  $\mathbb{G}_n \subset \mathbb{G}^{2^{n-1}} \subset \mathbb{G}_{(n-1)\log(2)/\log(p)}$ .*

*Proof.* — That  $a_n \leq p^n$  follows from Proposition 4.4 and the fact that  $\mathbb{G}_n/\mathbb{G}_{n+1}$  is elementary  $p$ -abelian. The second part follows from the first. □

### 4.1.2. The Golod–Shafarevich polynomial

Consider a minimal presentation of a finitely generated pro- $p$  group  $\mathbb{G}$ :

$$1 \longrightarrow \mathbb{R} \longrightarrow \mathbb{F} \xrightarrow{\varphi} \mathbb{G} \longrightarrow 1.$$

Suppose that  $\mathbb{R}/\mathbb{R}^p[\mathbb{R}, \mathbb{R}]$  is generated as an  $\mathbb{F}_p[[\mathbb{F}]]$ -module by the family  $\mathcal{F} = (\rho_i)$  of elements  $\rho_i \in \mathbb{F}$ . For  $k \geq 2$ , put  $r_k = |\{\rho_i, \omega(\rho_i) = k\}|$ ; here we assume the  $r_i$ 's finite.

DEFINITION 4.6. — *The series  $P(t) = 1 - dt + \sum_{k \geq 2} r_k t^k$  is a Golod–Shafarevich series associated to the presentation  $\mathcal{F}$  of  $\mathbb{G}$ .*

The theorem of Golod–Shafarevich asserts the following: if for some  $t_0 \in (0, 1)$  one has  $P(t_0) = 0$ , then  $\mathbb{G}$  is infinite (see [1] or [38]). Observe that when no information on the depth of the  $\rho_i$  is available, then one may take  $1 - dt + rt^2$  as Golod–Shafarevich series for  $\mathbb{G}$ , where  $r = d(H^2(\mathbb{G}))$ .

*Remark 4.7.* — When  $G = G_\emptyset$ , the  $p$ -rank of  $G_n$  corresponds to the  $p$ -rank of the class group of  $K_n$ , where  $K_n = K_\emptyset^{G_n}$ . Hence by Class Field Theory and with the help of a software package, in a certain sense it is easier to test if an element of  $G$  is in  $G_n$  than if it is in  $G^n$ .

## 4.2. Minkowski units and the Golod–Shafarevich polynomial of $G_\emptyset$

### 4.2.1. The principle

Let  $S$  be a finite saturated set of tame places of  $K$  as in Lemma 1.5, i.e. such that  $H^1(G_\emptyset) \simeq H^1(G_S)$  and  $|S| = d(V_{K,\emptyset})$ . Put  $d = d(G_\emptyset)$ . Let  $F$  be the free pro- $p$  group on  $d$  generators  $x_1, \dots, x_d$ . Consider now the minimal presentations of  $G_\emptyset$  and  $G_S$  induced by  $F$ , and the following diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & R_S & \xrightarrow{i'} & F & \xrightarrow{\varphi_S} & G_S \longrightarrow 1 \\
 & & \downarrow & & \downarrow = & & \downarrow \cong \\
 1 & \longrightarrow & R_\emptyset & \xrightarrow{i} & F & \xrightarrow{\varphi} & G_\emptyset \longrightarrow 1
 \end{array}$$

Put  $H_S = \ker(G_S \rightarrow G_\emptyset)$ ; the pro- $p$  group  $H_S$  is the normal subgroup of  $G_S$  generated by the tame inertia groups  $\langle \tau_p \rangle_{p \in S}$ . Hence, this diagram induces the following exact sequence

$$1 \longrightarrow R_S \longrightarrow R_\emptyset \xrightarrow{\psi'} H_S \longrightarrow 1,$$

where  $\psi' = \varphi_S \circ i$ . The Hochschild–Serre spectral sequences induce the following isomorphisms

$$\begin{array}{ccc}
 & H^2(G_\emptyset)^\wedge & \\
 \swarrow \cong & & \searrow \cong \\
 R_\emptyset/R_\emptyset^p[R_\emptyset, F] & \xrightarrow[\cong]{\psi} & H_S/H_S^p[H_S, G_\emptyset]
 \end{array}$$

where  $\psi$  is induced by  $\psi'$ . Using  $\psi$  we will study the depth of the relations of  $G_\emptyset$ : indeed,  $R_\emptyset/R_\emptyset^p[R_\emptyset, F]$  and  $H_S/H_S^p[H_S, G_\emptyset]$  inherit the Zassenhaus valuation from  $R_\emptyset$  and  $H_S$ , and thus the Zassenhaus valuation of  $F$ . Therefore an element of depth  $k$  in  $R_\emptyset/R_\emptyset^p[R_\emptyset, F]$  corresponds to an element of depth  $k$  in  $H_S/H_S^p[H_S, G_\emptyset]$ .

4.2.2. Minkowski elements

Here we extend the notion of Minkowski unit to the notion of Minkowski element. Set  $\mathcal{V}_K = V_{K,\emptyset}/(K^\times)^p$ .

DEFINITION 4.8. — *Let  $L/K$  be a Galois extension with Galois group  $G$ . We denote by  $\lambda'_{L/K} := t_G(\mathcal{V}_L)$  the  $\mathbb{F}_p[G]$ -rank of  $\mathcal{V}_L$ . One says that  $L/K$  has a Minkowski element if  $\lambda'_{L/K} \geq 1$ .*

LEMMA 4.9. — *One has  $\lambda'_{L/K} \geq \lambda_{L/K}$ , so the existence of a Minkowski unit implies that of a Minkowski element.*

*Proof.* — This follows immediately from the exact sequence

$$(4.1) \quad 1 \longrightarrow \mathcal{E}_L \longrightarrow \mathcal{V}_L \longrightarrow \text{Cl}_L[p] \longrightarrow 1. \quad \square$$

When  $L/K$  is a subextension of  $K_\emptyset/K$  one may give an upper bound for  $\lambda'_{L/K}$ :

PROPOSITION 4.10. — *Let  $L/K$  be a nontrivial finite Galois extension in  $K_\emptyset/K$ . Then  $\lambda'_{L/K} \leq d - 1 + r_1 + r_2$ . Moreover, if  $K_\emptyset/K$  is infinite then there exist infinitely many Galois extensions  $L/K$  in  $K_\emptyset/K$  such that  $\lambda'_{L/K} < d - 1 + r_1 + r_2$ .*

*Proof.* — Set  $H = \text{Gal}(K_\emptyset/L)$ . By Schreier's inequality (see [27], Corollary 3.6.3),  $d(\text{Cl}_L) = d(H) \leq |G/H|(d - 1) + 1$ . Hence by (4.1), we get

$$d(\mathcal{V}_L) \leq |G/H|(d - 1 + r_1 + r_2) + \delta,$$

showing that  $\lambda'_{L/K} \leq d - 1 + r_1 + r_2$ .

Suppose now that  $G_\emptyset$  is infinite and, except for finitely many Galois extensions  $L/K$  in  $K_\emptyset/K$ , one has  $\lambda'_{L/K} = d - 1 + r_1 + r_2$ . Then

$$\begin{aligned} d(\mathcal{V}_L) &\geq |G/H|(d - 1 + r_1 + r_2) \\ \text{and } d(\text{Cl}_L) &\geq 1 - \delta + |G/H|(d - 1) \geq |G/H|(d - 1), \end{aligned}$$

implying

$$-\chi_1(H) + 1 \geq -|G/H|\chi_1(G_\emptyset).$$

By [29, Proposition 1], the Galois group  $G_\emptyset$  must be free pro- $p$ , which is impossible. □

The converse below of Lemma 1.11 follows easily from the Chebotarev density theorem:

PROPOSITION 4.11. — *Let  $L/K$  be a finite  $p$ -extension with Galois group  $G$ .*

- (i) *If  $t_G(\mathcal{E}_{L,\emptyset}) \geq k$ , then there exist infinitely many sets  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_k\}$  consisting of  $k$  distinct tame primes of  $K$  such that  $\#G_{L,S}^{ab} = \#G_{L,\emptyset}^{ab}$ .*
- (ii) *If  $t_G(\mathcal{V}_L) \geq k$ , then there exist infinitely many sets  $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_k\}$  consisting of  $k$  distinct tame primes of  $K$  such that  $d(G_{L,S}) = d(G_{L,\emptyset})$ .*

From the computational view point, we will now consider the sequence of fields  $(K_n)$  in  $K_\emptyset/K$  induced by the Frattini filtration  $(G_n)$ : in other word,  $K_n = K_\emptyset^{G_n}$ . Put  $H_n = \text{Gal}(K_n/K)$ , and denote by  $\lambda'_n := \lambda'_{K_n}$  the  $\mathbb{F}_p[H_n]$ -free rank of  $\mathcal{V}_{K_n}$ .

Put  $d := d(G_\emptyset)$ , and  $r_{\max} := d + d(\mathcal{O}_K^\times)$ .

THEOREM 4.12. — *Take  $n \geq 2$ . Then  $G_\emptyset$  can be generated by  $d(G_\emptyset)$  generators and  $r_{\max}$  relations  $\{\rho_1, \dots, \rho_{r_{\max}}\}$  such that at least  $\lambda'_n$  relations are of depth greater than  $2^n$ .*

*Proof.* — We are assuming that the  $\mathbb{F}_p[H_n]$ -module  $\mathcal{V}_{K_n}$  is isomorphic to  $\mathbb{F}_p[H_n]^{\lambda'_n} \oplus N$  where  $N$  is torsion. Using Chebotarev’s theorem, choose a set  $S' = \{\mathfrak{p}_1, \dots, \mathfrak{p}_{\lambda'_n}\}$  of primes of  $K$  such that

- Each  $\mathfrak{p}_i$  splits completely from  $K$  to  $K_n$ ,
- The Frobenius at a prime  $\mathfrak{P}_{ij}$  of  $K_n$  above  $\mathfrak{p}_i$  in  $\text{Gal}(K'_n(\sqrt[p]{\mathcal{V}_{K_n}})/K'_n)$  lies in the  $i$ th copy of  $\mathbb{F}_p[H_n] \subset \mathcal{V}_{K_n}$  and generates that copy of  $\mathbb{F}_p[H_n]$  under the action of  $H_n$ .

We claim  $d(G_{K_n,S'}) = d(G_{K_n,\emptyset})$ . Indeed, there are  $|H_n|$  primes  $\mathfrak{P}_{ij}$  of  $K_n$  above  $\mathfrak{p}_i$  and they have independent Frobenius automorphisms in  $\text{Gal}(K'_n(\sqrt[p]{\mathcal{V}_{K_n}})/K'_n)$  by choice, even as we take the union over  $i$  from 1 to  $\lambda'_n$ . Gras–Munnier (Theorem 1.1) gives the equality. In fact, it gives more:  $d(G_{K_m,S'}) = d(G_{K_m,\emptyset})$  for all  $m < n$ . If this were false for  $m_0 < n$ , there would exist a  $\mathbb{Z}/p$ -extension of  $L/K_{m_0}$  ramified at primes (above those) of  $S'$ . Thus  $LK_n/K_n$  would be a  $\mathbb{Z}/p$ -extension ramified only at primes (above those) of  $S'$  contradicting the result for  $n$ . We have shown that the  $p$ -Frattini towers of  $G_{S'}$  and  $G_\emptyset$  agree at the first  $n$  levels. Thus the generators  $\tau_{\mathfrak{p}_i}$  of the tame inertia groups all have depth  $2^n$  in  $G_{S'}$ .

We have

$$0 \longrightarrow \text{III}_{S'}^2 \longrightarrow H^2(G_{S'}) \xrightarrow{res} \bigoplus_{\mathfrak{p}_i \in S'} H^2(G_{\mathfrak{p}_i})$$

and  $\dim \text{III}_{S'}^2 \leq \dim \mathbb{E}_{S'} = r_{\max} - \lambda'_n$ . We can say nothing about the depth of the relations coming from  $\text{III}_{S'}^2$ , so we assume they have minimal depth

two. The local relations are of the form  $[\sigma_{\mathfrak{p}_i}, \tau_{\mathfrak{p}_i}]_{\tau_{\mathfrak{p}_i}}^{1-N(\mathfrak{p}_i)}$  and are easily seen to have depth at least  $2^n + 1$ . As

$$G_{S'} / \langle \tau_{\mathfrak{p}_1}, \dots, \tau_{\mathfrak{p}_{\lambda'_n}} \rangle \simeq G_\emptyset$$

and taking this quotient trivializes the local relations, the theorem follows.  $\square$

**COROLLARY 4.13.** — *Assuming the hypothesis of Theorem 4.12, we may take  $1 - dt + (r_{\max} - \lambda'_n)t^2 + \lambda'_n t^{2^n}$  as a Golod–Shafarevich polynomial for  $G_\emptyset$ .*

*Example 4.14.* — Let us return to the field  $K = \mathbb{Q}(\sqrt{5 \cdot 13 \cdot 17 \cdot 19})$  of Section 1.3.3. Take  $H = K_2$ ,  $G = \text{Gal}(H/K)$ . As seen earlier,  $t_G(\mathcal{V}_H) \geq 1$ . Indeed, the existence of a Minkowski element follows from that of a Minkowski unit. Here a Golod–Shafarevich polynomial of  $G_\emptyset$  can be taken to be  $1 - 3t + 4t^2 + t^4$  instead of the naive choice  $1 - 3t + 5t^2$ .

### 4.2.3. The converse

Theorem 4.12 shows that the presence of Minkowski elements in the tower implies the existence of very deep relations in  $G_\emptyset$ . Here we show the converse, that the existence of very deep relations implies the presence of Minkowski elements.

For  $n \geq 1$ , recall that  $F$  is a free pro- $p$  group on  $d$  generators,  $F^m$  and  $F_m$  are the Zassenhaus and Frattini filtrations, and  $a_n$  is the smallest integer such that  $F^{a_n} \subset F_{n+1}$ . Recall from Lemma 4.1 that  $F_n \subset F^{2^{n-1}}$ . See Section 4.1.1. Put  $H_n = G_\emptyset/G_n$  and  $K_n = K_\emptyset^{G_n}$ .

**THEOREM 4.15.** — *Suppose that all the relations of  $G_\emptyset$  are of depth at least  $a_n$ . Then*

- (i) if  $\zeta_p \in K$ ,  $\lambda'_{K_n/K} \geq r_1 + r_2$ ;
- (ii) if  $\zeta_p \notin K$ ,  $\lambda'_{K_n/K} = r_1 + r_2 - 1 + d$ .

*Proof.* — Since all the relations of  $G_\emptyset$  have depth  $a_n$ , we see that  $G_\emptyset/G_\emptyset^{a_n} \simeq F/F^{a_n}$  has maximal Zassenhaus filtration for the first  $a_n$  steps. Thus for any set  $S$  satisfying  $d(G_S) = d(G_\emptyset)$  we have

$$F/F^{a_n} \simeq G_\emptyset/G_\emptyset^{a_n} \simeq G_S/G_S^{a_n}$$


and since  $F^{a_n} \subset F_{n+1}$ , we also have

$$F/F_{n+1} \simeq G_\emptyset/G_{\emptyset, n+1} \simeq G_S/G_{S, n+1}$$

so all relations of  $G_\emptyset$  have depth at least  $n + 1$  in the Frattini filtration.



We first address the case  $\zeta_p \in K$ . Consider the  $p$ -elementary abelian extensions  $K(\sqrt[p]{V_{K,\theta}})/K$  and  $K_2/K$ , the latter being the maximal unramified  $p$ -elementary abelian extension of  $K$ . By Kummer theory each is formed by adjoining to  $K$  the  $p$ th roots of elements  $\alpha \in K$ . Since  $K_2/K$  is everywhere unramified,  $(\alpha)$  is the  $p$ th power of an ideal, that is  $\alpha \in V_{K,\theta}$  so  $K(\sqrt[p]{V_{K,\theta}}) \supset K_2$  and  $d(\text{Gal}(K(\sqrt[p]{V_{K,\theta}})/K_2)) = r_1 + r_2$ . Note  $K_n \cap K(\sqrt[p]{V_{K,\theta}}) = K_2$  as the intersection is both unramified over  $K$  and  $p$ -elementary abelian over  $K$ . Let  $S := \{\mathfrak{p}_1, \dots, \mathfrak{p}_{r_1+r_2}\}$  consist of primes that split completely from  $K$  to  $K_2$  to  $K_n$  and whose Frobenius automorphisms form a basis of  $\text{Gal}(K(\sqrt[p]{V_{K,\theta}})/K_2)$ .




By the above discussion

$$(4.2) \quad F/F_{n+1} \simeq G_\theta/G_{\theta,n+1} \simeq G_S/G_{S,n+1}.$$

This will imply  $\lambda'_{K_n/K} \geq r_1 + r_2$ . Indeed, above each  $\mathfrak{p}_i$  there are  $[K_n : K]$  primes  $\mathfrak{P}_{ij}$  in  $K_n$  upon which  $\text{Gal}(K_n/K)$  acts transitively. If for some  $i$  the Frobenius automorphisms of the  $\mathfrak{P}_{ij}$  did not generate a distinct copy of  $\mathbb{F}_p[G_n]$  in  $\text{Gal}(K_n(\sqrt[p]{V_{K_n,\theta}})/K_n)$ , then there would be a dependence relation among them and by Gras–Munnier we would have  $d(G_{K_n,S}) > d(G_{K_n,\theta})$ , contradicting (4.2). Thus  $\lambda'_n \geq r_1 + r_2$  completing the proof in the  $\delta = 1$  case.

We now consider the case  $\zeta_p \notin K$ . As usual, the key fact is that  $K'_n \cap K'(\sqrt[p]{V_{K,\theta}}) = K'$  (following the proof of Proposition 2.8) so  $d(\text{Gal}(K'_n(\sqrt[p]{V_{K,\theta}})/K'_n)) = r_1 + r_2 - 1 + d$ .



We choose  $S := \{\mathfrak{p}_1, \dots, \mathfrak{p}_{r_1+r_2-1+d}\}$  to consist of primes of  $K$  that split completely from  $K$  to  $K'$  to  $K'_n$  and whose Frobenius automorphisms form a basis of  $\text{Gal}(K(\sqrt[r]{V_{K,\emptyset}})/K')$ . We complete the proof exactly as in the  $\zeta_p \in K$  case.  $\square$

**COROLLARY 4.16.** — *If all the relations of  $G_\emptyset$  are of depth at least  $p^2$  then  $K_2$  has a Minkowski element.*

*Proof.* — This follows immediately from Proposition 4.5 and Theorem 4.15.  $\square$

### 4.3. Theorem 2.9 revisited

There is another way by which we can obtain Theorem 2.9 in the context of Golod–Shafarevich series  $P(t)$ . Indeed, such a series for a pro- $p$  group  $G$  approximates the Hilbert series  $H_G(t)$  of the Zassenhaus filtration of  $G$ . In particular the Golod–Shafarevich Theorem is a consequence of this inequality: if there is some  $t_0 \in ]0, 1[$  such that  $P(t_0) < 0$  then necessarily  $H_G(t_0)$  diverges, implying the infiniteness of  $G$ .

Retain the notations of Section 2.3, and fix  $n \gg 0$ . Apply Corollary 4.13 to  $K_n/K$  by taking  $1 - dt + (r_{\max} - \lambda)t^2 + \lambda t^{2n}$  as a Golod–Shafarevich polynomial for  $G_\emptyset$ . Now, as  $n$  can be arbitrarily large, we see that  $1 - dt + (r_{\max} - \lambda)t^2$  is a Golod–Shafarevich polynomial for  $G_\emptyset$ .

Of course, the question of determining  $\lambda$  when it is nonzero seems a hard problem, except in the case where at the beginning of the tower, we see  $\lambda = 0$ . Here is an explicit alternative.

**COROLLARY 4.17.** — *Let  $n \in \mathbb{Z}_{>1}$ . One has:*

- (i) *if  $t_{H_n}(\mathcal{E}_{K_n}) = 0$  and  $\beta = 0$ , then  $\text{Def}(G_\emptyset) = r_1 + r_2 - 1 + \delta$ ;*
- (ii) *if  $t_{H_n}(\mathcal{E}_{K_n}) = \lambda_n > 0$ , then one may take  $1 - dt + (r_{\max} - \lambda_n)t^2 + \lambda_n t^{2n}$  as a Golod–Shafarevich polynomial for  $G_\emptyset$ .*

**Remark 4.18.** — The condition  $\beta = 0$  can be relaxed as noted in Theorem 3.14.

## 5. The case of imaginary quadratic fields

In this section, we take  $p = 2$  and let  $K := \mathbb{Q}(\sqrt{D})$  be an imaginary quadratic field of discriminant  $D < -7$ . Since the unit rank of  $K$  is 1, we have  $\text{Def}(G_\emptyset) \in \{0, 1\}$ . In this simplest of all non-trivial situations, we will discuss the deficiency of  $G_\emptyset$  and explore the extent to which we can detect relations using the machinery and notation set up in Section 2.2.

### 5.1. The frame

Let  $d = d(\text{Cl}_K)$  be the 2-rank of the class group of  $K := \mathbb{Q}(\sqrt{D})$ . By Gauss's genus theory, we know that  $D$  admits a unique (up to reordering) factorization into  $d + 1$  integers, each of which is a "prime fundamental discriminant" – meaning it is the discriminant of a quadratic field in which a single prime ramifies. For an odd prime  $q$ , we define  $q^* := (-1)^{(q-1)/2}q$ . The prime discriminants are then  $q^*$  as  $q$  ranges over all odd primes, as well as  $-4$  and  $\pm 8$ . We write  $D = q_1^* \cdots q_{d+1}^*$ , with the convention that if  $D$  is even, then  $q_{d+1}^* \in \{-4, -8, 8\}$ .

Put  $q_0^* = -1$  and for each  $i$  in the range  $0 \leq i \leq d$ , put

$$K_i := K(\sqrt{q_0^*}, \dots, \sqrt{q_{i-1}^*}, \sqrt{q_{i+1}^*}, \dots, \sqrt{q_d^*}),$$

where

$$q'_d = \begin{cases} q_d^* & \text{if } D \text{ is odd} \\ q_d^* & \text{if } q_{d+1}^* = \pm 8 \\ 2 & \text{if } q_{d+1}^* = -4. \end{cases}$$

Also define  $L' := K(\sqrt{q_0^*}, \sqrt{q_1^*}, \dots, \sqrt{q_{d-1}^*}, \sqrt{q'_d})$ . A direct computation shows that the number field  $L'$  is the governing field  $K(\sqrt{V_\emptyset})$  (see Section 2.2). Choose prime numbers  $p_0, \dots, p_d$  that split in  $K$  and such that for each  $i$  in the range  $0 \leq i \leq d$ , the Frobenius automorphisms of the  $p_j, j \neq i$  in  $L'/\mathbb{Q}$  generate the Galois group of the quadratic extension  $L'/K_i$ . Fix a prime  $\mathfrak{p}_i | p_i$  of  $K$  and put  $S_2 = \{\mathfrak{p}_0\}$ ,  $S_1 = \{\mathfrak{p}_1, \dots, \mathfrak{p}_d\}$ , and  $S = S_1 \cup S_2$ . Observe that the primes  $p_1, \dots, p_d$  all are congruent to 1 mod 4 and that  $p_0 \equiv 3 \pmod{4}$ .

As the 2-part of the class group of  $K$  has  $d$  generators, Lemma 2.5 shows the existence of  $d$  independent quadratic extensions  $F_i$  above  $K_\emptyset$ , totally ramified at  $\mathfrak{p}_i$ ,  $i = 1, \dots, d$ , so  $d(X_S) \geq d$ . This puts us in the situation where where a difficult relation, if it exists, is detectable by the set  $S_2$ . Now, by studying the Galois module structure of units in imaginary biquadratic number fields, we can specify conditions under which  $\text{Def}(G_\emptyset) = 1$ ; see Theorem 5.3 below.

LEMMA 5.1. — *Let  $K_0/\mathbb{Q}$  be a real quadratic field;  $G_0 = \text{Gal}(K_0/\mathbb{Q})$ . Then  $\mathcal{E}_{K_0}$  is  $\mathbb{F}_2[G_0]$ -free if and only if, the norm of the fundamental unit  $\varepsilon$  is  $-1$ . More precisely, as an  $\mathbb{F}_2[G_0]$ -module,*

$$\mathcal{E}_K \simeq \begin{cases} \mathbb{F}_2 \oplus \mathbb{F}_2 & N(\varepsilon) = 1 \\ \mathbb{F}_2[G_0] & N(\varepsilon) = -1. \end{cases}$$

*Proof.* — If the norm of  $\varepsilon$  is  $+1$ , then modulo  $(\mathcal{O}_K^\times)^2$ , we get  $\varepsilon^\sigma \equiv \varepsilon$ . If the norm of  $\varepsilon$  is  $-1$ , then  $\mathcal{E}_K$  is generated by  $\varepsilon(\mathcal{O}_K^\times)^2$  as  $G$ -module, and  $\langle \varepsilon(\mathcal{O}_K^\times)^2 \rangle$  is  $\mathbb{F}_2[G_0]$ -free.  $\square$

Recall this well-known result:

LEMMA 5.2. — *Let  $F/\mathbb{Q}$  be an imaginary biquadratic field. Let  $K_0$  be the real quadratic subfield, and let  $\varepsilon$  be the fundamental unit of  $K_0$ . Then,  $|\mathcal{O}_F^\times / \langle \mu_F, \varepsilon \rangle| = 1$  or  $2$ . In particular, if  $F/K_0$  is ramified at some odd prime, then  $\mathcal{O}_F^\times = \langle \mu_F, \varepsilon \rangle$ .*

### 5.2. Main result

We can now prove:

THEOREM 5.3. — *Let  $K$  be an imaginary quadratic field of discriminant  $D$ . Assume that we can write  $D = D_1 D_2$ , where  $D_1 > 0$  and  $D_2$  are fundamental discriminants, such that:*

- (i) *the norm of the fundamental unit of  $\mathbb{Q}(\sqrt{D_1})$  is  $+1$ ,*
- (ii) *some odd prime number divides  $D_2$ .*

*Then  $\text{Def}(G_\theta) = 1$ , and the difficult relation is detected by the quadratic extension  $K(\sqrt{D_1})/K$ .*

*Proof.* — Put  $F := K(\sqrt{D_1})$ . As  $D_1$  and  $D_2$  are fundamental discriminants, then  $F/K$  is unramified. By assumption (ii) and Lemma 5.2,  $\mathcal{O}_F^\times = \langle \varepsilon, -1 \rangle$ , where  $\varepsilon$  is the fundamental unit of  $\mathbb{Q}(\sqrt{D_1})$ . By assumption (i) and Lemma 5.1,  $\mathcal{E}_F$  is not  $\mathbb{F}_2[G]$ -free, where  $G = \text{Gal}(F/K)$ : in other words  $t_G(\mathcal{E}_F) = 0$ . The result follows by Theorem 3.14 (here  $\sqrt{-1} \notin F$ ).  $\square$

Remark 5.4. — To elaborate further, observe that  $p_0$  splits in  $F/K$ . Indeed, by the choice of  $p_0$  we have, for  $i = 1, \dots, d - 1$ ,  $\left(\frac{q_i^*}{p_0}\right) = \left(\frac{q'_i}{p_0}\right) = 1$ . Let us study two cases.

(a). — Suppose first that  $q'_d = q_d^*$ . Then by recalling that  $\left(\frac{D}{p_0}\right) = 1$ , one also gets  $\left(\frac{q_{d+1}^*}{p_0}\right) = 1$ , and then  $\left(\frac{D_1}{p_0}\right) = 1$  (in this case  $D_1$  is the product of some of the  $q_i^*$ ).

(b). — Suppose now that  $q'_d = 2$ . Since  $p_0 \equiv 3 \pmod{4}$  and  $D = q_1^* \dots q_{d+1}^*$ , we have  $\left(\frac{q_d^*}{p_0}\right) = -1$ . By assumption, there exists an odd prime  $p$  that divides  $D_2$ . We may choose  $p = q_d$  (before fixing  $p_0$ ). Then,  $D_1$  is the product of various  $q_i^*$ , for  $i = 1, \dots, d - 1$  so  $\left(\frac{D_1}{p_0}\right) = 1$ .

As  $p_0$  splits completely in  $F/K$ , we see  $\prod_{\mathfrak{q}|\mathfrak{p}_0} \mathcal{U}_{\mathfrak{q}}/\mathcal{U}_{\mathfrak{q}}^2$  is  $\mathbb{F}_2[G]$ -free of rank 1. But as  $t_G(\mathcal{E}_F) = 0$ , the subgroup  $I_{\mathfrak{p}_0}$  of  $\text{RCG}_F(\mathfrak{p}_0)$  generated by

the ramification at  $\mathfrak{p}_0$  is not trivial. Put  $I := I_{\mathfrak{p}_0}/I_{\mathfrak{p}_0}^2$ . By Nakayama's lemma, the coinvariants  $I_G$  are also not trivial, hence there exists at least one quadratic extension  $F_1/F_\theta$ , Galois over  $K$ , totally ramified at some  $\mathfrak{P}|\mathfrak{p}_0$ , such that  $G$  acts trivially on  $\text{Gal}(F_1/F_\theta)$ . The compositum  $F_1K_\theta/K_\theta$  is ramified at  $\mathfrak{p}_0$  and produces a  $(d+1)$ st relation. This is the formalism of Example 1.6.

**COROLLARY 5.5.** — *Let  $K$  be an imaginary quadratic field of discriminant  $D$ . Suppose  $D$  is divisible by at least two odd primes  $p_1, p_2$  such that  $p_1 \equiv p_2 \equiv 3 \pmod{4}$ . Then  $\text{Def}(G_\theta) = 1$ .*

*Proof.* — If there is another odd prime  $q$  that divides  $D$ , take  $D_1 = p_1 p_2$ . If  $K = \mathbb{Q}(\sqrt{-p_1 p_2})$  (resp.  $\mathbb{Q}(\sqrt{-2p_1 p_2})$ ), take  $D_1 = 4p_1$  (resp.  $D_1 = 8p_1$ ).  $\square$

*Example 5.6 (Martinet [22]).* — Take  $K = \mathbb{Q}(\sqrt{-21})$ . Then, by Odlyzko bounds the 2-tower  $K_\theta/K$  is finite, and it is not hard to see  $G_\theta \simeq (\mathbb{Z}/2\mathbb{Z})^2$ , and  $\text{Def}(G_\theta) = 1$ .

*Example 5.7 (See Example 1.6).* — Take  $K = \mathbb{Q}(\sqrt{-5460})$ ,  $D_1 = 21$  and  $D_2 = -260$ . We then get a difficult relation coming from the extension  $K(\sqrt{21})/K$ , and  $\text{Def}(G_\theta) = 1$ .

**COROLLARY 5.8.** — *Suppose  $k \geq 2$ , and  $p_1, \dots, p_k$  are  $k$  distinct odd primes, exactly one of which, say  $p_1$ , is  $\equiv 3 \pmod{4}$ . For the imaginary quadratic field  $K = \mathbb{Q}(\sqrt{-2p_1 \dots p_k})$  with discriminant  $D = -8p_1 \dots p_k$ , we have:  $\text{Def}(G_\theta) = 1$ .*

*Proof.* — Take  $D_1 = 8p_1$ .  $\square$

*Example 5.9.* — Take  $K = \mathbb{Q}(\sqrt{-p_1 p_2})$ , with primes  $p_1, p_2$  such that  $p_1 \equiv 1 \pmod{4}$  and  $p_2 \equiv 3 \pmod{4}$ . Here the hypotheses of Theorem 5.3 do not apply and  $r = d = 1$  so  $\text{Def}(G_\theta) = 0$ .

*Example 5.10.* — The hypotheses of Theorem 5.3 do not apply for  $K = \mathbb{Q}(\sqrt{-130})$ . As noted by Martinet [22], in that case,  $G_\theta$  is the quaternion group so  $r = d = 2$ .

*Example 5.11.* — Take  $K = \mathbb{Q}(\sqrt{-5 \cdot 13 \cdot 41})$ . Here  $r = d + 1 = 3$ ; indeed the norm of the fundamental unit of  $\mathbb{Q}(\sqrt{5 \cdot 41})$  is  $+1$ .

### 5.3. $\text{Def}(G_\theta)$ is maximal almost all the time

We easily deduce from Theorem 5.3 that the presence of a Minkowski unit in a quadratic unramified extension  $F/K$  is rare, with the consequence

that, generically, the deficiency of  $G_\emptyset$  is maximal. Let us say more precisely what we mean by the term “generically” here. Denote by  $\mathcal{F}$  the set of imaginary quadratic fields. For  $X \geq 2$ , put

$$\mathcal{F}(X) = \{K \in \mathcal{F}, |\text{disc}(K)| \leq X\},$$

and

$$\mathcal{F}_0(X) = \{K \in \mathcal{F}(X), \text{Def}(G_\emptyset) = 0\}.$$

**THEOREM 5.12.** — *There is an absolute constant  $C > 0$  such that for all  $X$  large enough,*

$$\frac{\#\mathcal{F}_0(X)}{\#\mathcal{F}(X)} \leq C \frac{\log \log X}{\sqrt{\log X}}.$$

*In particular, when ordered by absolute value of the discriminant, the proportion of imaginary quadratic fields for which  $\text{Def}(G_\emptyset) = 0$ , tends to zero when  $X \rightarrow \infty$ .*

*Proof.* — We use the analytic number theory tools of [21, Theorem 4.6] due to Fouvry. Let  $K$  be an imaginary quadratic field. Put

$$B(X) = \left\{ K \in \mathcal{F}(X), \exists 2 \text{ distinct odd primes } p \equiv q \equiv 3 \pmod{4}, \right. \\ \left. pq \mid \text{disc}(K) \right\}.$$

By Corollary 5.5, for every  $K \in B(X)$  one has  $\text{Def}(G_\emptyset) = 1$ . Hence  $\mathcal{F}_0(X)$  is in the complement  $C(X)$  of  $B(X)$ .

Denote by  $A_i(X)$  the set of square-free integers  $n \leq X$  having exactly  $i$  prime factors  $\equiv 3 \pmod{4}$ , put  $A(X) = A_0(X) \cup A_1(X)$ . Clearly,  $|C(X)| = O(|A(X)|)$ .

In the proof of Theorem 4.6 of [21], it is shown that uniformly for  $X$  large enough, one has  $|A_0(X)| = O(X/\sqrt{\log X})$  and  $|A_1(X)| = O\left(X \frac{\log \log X}{\sqrt{\log X}}\right)$ .

Thus  $|C(X)| = O\left(X \frac{\log \log X}{\sqrt{\log X}}\right)$ . We conclude by noting that  $|\mathcal{F}(X)| = \frac{3}{\pi^2} X + O(\sqrt{\log X})$  (see for example [8, Section 4]). □

The referee asked whether the bulk of the deficiency zero cases for  $p = 2$  and imaginary quadratic fields arise with tower group the quaternion group  $Q_8$  of order 8, suggesting a criterion from Table II of [2] as a possible method of proving this. Let  $\mathcal{F}_{Q_8}(X)$  be the number quadratic imaginary fields having discriminant bounded in absolute value by  $X$  with 2-tower group  $Q_8$ . While it is not difficult to show, using the GRH versions of the effective Chebotarev Theorem, that

$$c \frac{(\log \log X)^2}{\log(X)} \leq \frac{\#\mathcal{F}_{Q_8}(X)}{\#\mathcal{F}(X)}$$

for some  $c > 0$  and  $X$  large enough, showing that  $Q_8$  towers are 100% of the deficiency zero cases for  $p = 2$  seems difficult.

### Acknowledgments

We thank the referee for thoroughly reading the paper, making a number of helpful suggestions and asking some interesting questions.

### BIBLIOGRAPHY

- [1] I. V. ANDOŽSKI, “On some classes of closed pro- $p$ -groups”, *Math. USSR, Izv.* **9** (1965), no. 4, p. 663-691.
- [2] E. BENJAMIN, F. LEMMERMEYER & C. SNYDER, “Imaginary quadratic fields  $k$  with cyclic  $Cl_2(k^1)$ ”, *J. Number Theory* **67** (1997), no. 2, p. 229-245.
- [3] W. BOSMA, J. CANNON & C. PLAYOUST, “The MAGMA algebra system. I. The user language”, *J. Symb. Comput.* **24** (1997), no. 3-4, p. 235-265.
- [4] N. BOSTON & J. WANG, “The 2-class tower of  $\mathbb{Q}(\sqrt{-5460})$ ”, in *Geometry, algebra, number theory, and their information technology applications*, Springer Proceedings in Mathematics & Statistics, vol. 251, Springer, 2018, p. 71-80.
- [5] C. W. CURTIS & I. REINER, *Representation theory of finite groups and associative algebras*, Pure and Applied Mathematics, vol. 11, John Wiley & Sons, 1988.
- [6] J. D. DIXON, M. P. F. DU SAUTOY, A. MANN & D. SEGAL, *Analytic pro- $p$  groups*, 2nd ed., London Mathematical Society Lecture Note Series, vol. 157, Cambridge University Press, 1999.
- [7] J.-M. FONTAINE & B. MAZUR, “Geometric Galois representations”, in *Elliptic curves, modular forms, and Fermat’s last theorem*, Series in Number Theory, vol. 1, International Press, 1995, p. 41-78.
- [8] É. FOUVRY & J. KLÜNERS, “On the 4-rank of the class groups of quadratic number fields”, *Invent. Math.* **167** (2007), no. 3, p. 455-513.
- [9] E. S. GOLOD & I. R. SHAFAREVICH, “On the class field tower”, *Izv. Akad. Nauk SSSR, Ser. Mat.* **28** (1964), p. 261-272.
- [10] G. GRAS, *Class Field Theory, From theory to practice*, corr. 2nd ed., Springer Monographs in Mathematics, Springer, 2005.
- [11] ———, “On the  $T$ -ramified,  $S$ -split  $p$ -class field towers over an extension of degree prime to  $p$ ”, *J. Number Theory* **129** (2009), no. 11, p. 2843-2852.
- [12] G. GRAS & A. MUNNIER, “Extensions cycliques  $T$ -totalement ramifiées”, *Publ. Math. Besançon* **1998** (1998), article no. 6 (17 pages).
- [13] F. HAJIR, C. MAIRE & R. RAMAKRISHNA, “On the Shafarevich group of restricted ramification extensions of number fields in the tame case”, *Indiana Univ. Math. J.* **70** (2021), no. 6, p. 2693-2710.
- [14] H. KISILEVSKY, “Number Fields with Class Number congruent to 4 mod 8 and Hilbert’s Theorem 94”, *J. Number Theory* **8** (1976), p. 271-279.
- [15] H. KOCH, *Galois Theory of  $p$ -Extensions*, Springer Monographs in Mathematics, Springer, 2002.
- [16] J. LABUTE, “Mild pro- $p$ -groups and Galois groups of  $p$ -extensions of  $\mathbb{Q}$ ”, *J. Reine Angew. Math.* **596** (2006), p. 155-182.

- [17] J. LABUTE & J. MINÁČ, “Mild pro-2-groups and 2-extensions of  $\mathbb{Q}$  with restricted ramification”, *J. Algebra* **332** (2011), p. 136-158.
- [18] M. LAZARD, “Groupes analytiques  $p$ -adiques”, *Publ. Math., Inst. Hautes Étud. Sci.* **26** (1965), p. 389-603.
- [19] Y. LIU, M. M. WOOD & D. ZUREICK-BROWN, “A predicted distribution for Galois groups of maximal unramified extensions”, *Invent. Math.* **237** (2024), p. 49-116.
- [20] C. MAIRE, “Cohomology of number fields and analytic pro- $p$ -groups”, *Mosc. Math. J.* **10** (2010), no. 2, p. 399-414.
- [21] ———, “On the quotients of the maximal unramified 2-extensions of a number field”, *Doc. Math.* **23** (2018), p. 1263-1290.
- [22] J. MARTINET, “Tours de corps de classes et estimations de discriminants”, *Invent. Math.* **44** (1978), p. 65-73.
- [23] B. MAZUR, “Notes on étale cohomology of number fields”, *Ann. Sci. Éc. Norm. Supér.* **6** (1973), p. 521-553.
- [24] J. NEUKIRCH, A. SCHMIDT & K. WINGBERG, *Cohomology of Number Fields*, Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer, 2013, second edition, corrected second printing.
- [25] M. OZAKI, “Construction of maximal unramified  $p$ -extensions with prescribed Galois groups”, *Invent. Math.* **183** (2011), no. 3, p. 649-680.
- [26] THE PARI GROUP, “PARI/GP version 2.9.4”, 2018, available from <http://pari.math.u-bordeaux.fr/>.
- [27] L. RIBES & P. ZALESKII, *Profinite Groups*, 2nd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, vol. 40, Springer, 2010.
- [28] P. ROQUETTE, “On Class Field Towers”, in *Algebraic Number Theory* (J. W. S. Cassels & A. Fröhlich, eds.), Academic Press Inc., 1967, p. 231-249.
- [29] A. SCHMIDT, “Bounded defect in partial Euler characteristics”, *Bull. Lond. Math. Soc.* **28** (1996), no. 5, p. 463-464.
- [30] ———, “Rings of integers of type  $K(\pi, 1)$ ”, *Doc. Math.* **12** (2007), p. 441-471.
- [31] ———, “Über Pro- $p$ -Fundamentalgruppen markierter arithmetischer Kurven”, *J. Reine Angew. Math.* **640** (2010), p. 203-235.
- [32] A. SCHOLZ, “Über die Beziehung der Klassenzahlen quadratischer Körper zueinander”, *J. Reine Angew. Math.* **166** (1932), p. 201-203.
- [33] R. SCHOOF, “Infinite class field towers of quadratic fields”, *J. Reine Angew. Math.* **372** (1986), p. 209-220.
- [34] W. R. SCOTT, *Group Theory*, Dover Publications, 1987.
- [35] J.-P. SERRE, *Cohomologie Galoisienne*, 5th ed., Lecture Notes in Mathematics, vol. 5, Springer, 1997.
- [36] I. R. SHAFAREVICH, “Algebraic number fields”, in *Proceedings of the International Congress of Mathematicians 1962*, Inst. Mittag-Leffler, 1963, p. 163-176.
- [37] ———, “Extensions with prescribed ramification points”, *Publ. Math., Inst. Hautes Étud. Sci.* **18** (1964), p. 71-95, in Russian; English translation *Amer. Math. Soc. Transl.*, **59**, p. 128-149.
- [38] È. B. VINBERG, “On a theorem concerning on infinite dimensionality of an associative algebra”, *Izv. Akad. Nauk SSSR, Ser. Mat.* **29** (1965), p. 208-214, in Russian; English translation *Amer. Math. Soc. Transl. (2)* **82**, p. 237-242.
- [39] K. WINGBERG, “On Demuškin groups with involution”, *Ann. Sci. Éc. Norm. Supér.* **22** (1989), no. 4, p. 555-567.
- [40] ———, “On the Fontaine–Mazur conjecture for CM-fields”, *Compos. Math.* **131** (2002), no. 3, p. 341-354.
- [41] ———, “Free quotients of Demuškin groups with operators”, preprint, 2004.



Manuscrit reçu le 24 juillet 2021,  
révisé le 23 mai 2022,  
accepté le 10 novembre 2022.

Farshid HAJIR  
Department of Mathematics & Statistics  
University of Massachusetts  
Amherst, MA 01003, USA  
hajir@math.umass.edu

Christian MAIRE  
FEMTO-ST Institute, Université Bourgogne  
Franche-Comté, CNRS  
15B avenue des Montboucons  
25000 Besançon, France  
christian.maire@univ-fcomte.fr

Ravi RAMAKRISHNA  
Department of Mathematics, Cornell University  
Ithaca, USA  
ravi@math.cornell.edu